

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 May 2001 (31.05.2001)

PCT

(10) International Publication Number
WO 01/37779 A2

- (51) International Patent Classification: **A61K** (US). **RECIPON, Herve**; 85 Fortuna Avenue, San Francisco, CA 94115 (US). **SUN, Yongming**; Apartment 260, 869 S. Winchester Boulevard, Apartment 260, San Jose, CA 95128 (US).
- (21) International Application Number: PCT/US00/32056
- (22) International Filing Date: 22 November 2000 (22.11.2000) (74) Agents: **LICATA, Jane, Massey et al.**; Licata & Tyrrell P.C., 66 E. Main Street, Marlton, NJ 08053 (US).
- (25) Filing Language: English (81) Designated States (*national*): CA, JP.
- (26) Publication Language: English (84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).
- (30) Priority Data: 60/166,973 23 November 1999 (23.11.1999) US
- (71) Applicant: **DIADEXUS, INC.** [US/US]; 3303 Octavius Drive, Santa Clara, CA 95054 (US). **Published:**
— Without international search report and to be republished upon receipt of that report.
- (72) Inventors: **SALCEDA, Susana**; 4118 Crescendo Avenue, San Jose, CA 95136 (US). **CAFFERKEY, Robert**; Apartment 218, 350 Elan Village Lane, San Jose, CA 95134 For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/37779 A2

(54) Title: A NOVEL METHOD OF DIAGNOSING, MONITORING, STAGING, IMAGING AND TREATING BREAST CANCER

(57) Abstract: The present invention provides new markers and methods for detecting, diagnosing, monitoring, staging, prognosticating, imaging and treating breast cancer.

- 1 -

A NOVEL METHOD OF DIAGNOSING,
MONITORING, STAGING, IMAGING AND TREATING BREAST CANCER

FIELD OF THE INVENTION

This invention relates, in part, to newly identified
5 breast cancer specific genes and assays for detecting,
diagnosing, monitoring, staging, prognosticating, imaging and
treating cancers, particularly breast cancer.

BACKGROUND OF THE INVENTION

10 It is estimated that one out of every nine women in
America will develop breast cancer sometime during her life
based on a lifespan of 85 years. Annually, over 180,000 women
in the United States are diagnosed with breast cancer and
approximately 46,000 die from this disease. Every woman is
15 at risk for breast cancer. However, a woman's chances of
developing breast cancer increase as she grows older; 80
percent of all cancers are found in women over the age of 50.
There are also several risk factors that can increase a
woman's chances of developing breast cancer. These include
20 a family history of breast cancer, having no children or the
first child after the age of 30, and an early start of
menstruation. However, more than 70 percent of women who
develop breast cancer have no known risk factors. Less than
10 percent of breast cancer cases are thought to be related
25 to the BRCA1 gene discovered in 1994. Researchers are now
investigating the role of other factors such as nutrition,
alcohol, exercise, smoking, and oral contraceptives in
development of this gynecologic cancer. Mammograms, or
special x-rays of the breast, can detect more than 90 percent
30 of all cancers.

- 2 -

Procedures used for detecting, diagnosing, monitoring, staging, and prognosticating breast cancer are of critical importance to the outcome of the patient. Patients diagnosed early generally have a much greater five-year survival rate
5 as compared to the survival rate for patients diagnosed with distant metastasized breast cancer. New diagnostic methods which are more sensitive and specific for detecting early breast cancer are clearly needed.

Breast cancer patients are closely monitored following
10 initial therapy and during adjuvant therapy to determine response to therapy and to detect persistent or recurrent disease or metastasis. Thus, there is also clearly a need for cancer markers which are more sensitive and specific in detecting breast cancer recurrence.

15 Another important step in managing breast cancer is to determine the stage of the patient's disease. Stage determination has potential prognostic value and provides criteria for designing optimal therapy. Generally, pathological staging of cancer is preferable over clinical
20 staging because the former gives a more accurate prognosis. However, clinical staging would be preferred were it at least as accurate as pathological staging because it does not depend on an invasive procedure to obtain tissue for pathological evaluation. Staging of cancer would be improved by detecting
25 new markers in cells, tissues or bodily fluids which could differentiate between different stages of invasion.

New breast cancer specific genes, referred to herein as BCSGs, have now been identified for use in diagnosing, monitoring, staging, imaging and treating cancers, and in
30 particular breast cancer. Accordingly, the present invention relates to new methods for detecting, diagnosing, monitoring, staging, prognosticating, in vivo imaging and treating cancer via a BCSG. BCSG refers, among other things, to native proteins expressed by the genes comprising the polynucleotide
35 sequences of BCSG-1 or Gene ID 332369 (SEQ ID NO: 1), BCSG-2

- 3 -

or Gene ID 480489 (SEQ ID NO:2 or 18), BCSG-3 or Gene ID 274731 (SEQ ID NO:3 or 20), BCSG-4 or Gene ID 173388 (SEQ ID NO:4) or BCSG-5 or Clone ID 3040232, Gene ID 411152 (SEQ ID NO:5). Exemplary proteins expressed by genes BCSG-2 and BCSG-3 are depicted herein as SEQ ID NO:19 and SEQ ID NO:21. By "BCSG" it is also meant herein variant polynucleotides which, due to degeneracy in genetic coding, comprise variations in nucleotide sequence as compared to SEQ ID NO: 1, 2, 3, 4, 5, 18 or 20 but which still encode the same proteins. In the alternative, what is meant by BCSG as used herein, means the native mRNAs encoded by the genes comprising BCSG-1 or Gene ID 332369 (SEQ ID NO: 1), BCSG-2 or Gene ID 480489 (SEQ ID NO:2 or 18), BCSG-3 or Gene ID 274731 (SEQ ID NO:3 or 20), BCSG-4 or Gene ID 173388 (SEQ ID NO:4) or BCSG-5 or Clone ID 3040232, Gene ID 411152 (SEQ ID NO:5) or it can refer to the actual genes comprising BCSG-1 or Gene ID 332369 (SEQ ID NO: 1), BCSG-2 or Gene ID 480489 (SEQ ID NO:2 or 18), BCSG-3 or Gene ID 274731 (SEQ ID NO:3 or 20), BCSG-4 or Gene ID 173388 (SEQ ID NO:4) or BCSG-5 or Clone ID 3040232, Gene ID 411152 (SEQ ID NO:5), or levels of polynucleotides which are capable of hybridizing under stringent conditions to the antisense sequences of SEQ ID NO: 1, 2, 3, 4, 5, 18 or 20.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

- 4 -

SUMMARY OF THE INVENTION

Toward these ends, and others, it is an object of the present invention to provide BCSGs comprising a polynucleotide of SEQ ID NO:1, 2, 3, 4, 5, 18 or 20 or a variant thereof, a
5 protein expressed by a polynucleotide of SEQ ID NO:1, 2, 3, 4, 5, 18 or 20 or variant thereof which expresses the protein; or a polynucleotide which is capable of hybridizing under stringent conditions to the antisense sequence of SEQ ID NO: 1, 2, 3, 4, 5, 18 or 20.

10 Further provided is a method for diagnosing the presence of breast cancer by analyzing for changes in levels of BCSG in cells, tissues or bodily fluids compared with levels of BCSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein a change in levels of
15 BCSG in the patient versus the normal human control is associated with breast cancer.

Further provided is a method of diagnosing metastatic breast cancer in a patient having breast cancer which is not known to have metastasized by identifying a human patient
20 suspected of having breast cancer that has metastasized; analyzing a sample of cells, tissues, or bodily fluid from such patient for BCSG; comparing the BCSG levels in such cells, tissues, or bodily fluid with levels of BCSG in preferably the same cells, tissues, or bodily fluid type of
25 a normal human control, wherein an increase in BCSG levels in the patient versus the normal human control is associated with breast cancer which has metastasized.

Also provided by the invention is a method of staging breast cancer in a human by identifying a human patient having
30 breast cancer; analyzing a sample of cells, tissues, or bodily fluid from such patient for BCSG; comparing BCSG levels in such cells, tissues, or bodily fluid with levels of BCSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein an increase in BCSG levels in
35 the patient versus the normal human control is associated with

- 5 -

a cancer which is progressing and a decrease in the levels of BCSG is associated with a cancer which is regressing or in remission.

Further provided is a method of monitoring breast cancer in a human patient for the onset of metastasis. The method comprises identifying a human patient having breast cancer that is not known to have metastasized; periodically analyzing cells, tissues, or bodily fluid from such patient for BCSG; comparing the BCSG levels in such cells, tissue, or bodily fluid with levels of BCSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein an increase in BCSG levels in the patient versus the normal human control is associated with a cancer which has metastasized.

Further provided is a method of monitoring the change in stage of cancer in a human patient having breast cancer by looking at levels of BCSG in the human patient. The method comprises identifying a human patient having breast cancer; periodically analyzing cells, tissues, or bodily fluid from such patient for BCSG; comparing the BCSG levels in such cells, tissue, or bodily fluid with levels of BCSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein an increase in BCSG levels in the patient versus the normal human control is associated with breast cancer which is progressing and a decrease in the levels of BCSG is associated with breast cancer which is regressing or in remission.

Further provided are methods of designing new therapeutic agents targeted to BCSGs for use in imaging and treating cancer. For example, in one embodiment, therapeutic agents such as antibodies targeted against a BCSG or fragments of such antibodies can be used to treat, detect or image localization of a BCSG in a patient for the purpose of detecting or diagnosing a disease or condition. In this embodiment, an increase in the amount of labeled antibody

- 6 -

detected as compared to normal tissue would be indicative of tumor metastases or growth. Such antibodies can be polyclonal, monoclonal, or omniclonal or prepared by molecular biology techniques. The term "antibody", as used herein and
5 throughout the instant specification is also meant to include aptamers and single-stranded oligonucleotides such as those derived from an *in vitro* evolution protocol referred to as SELEX and well known to those skilled in the art. Antibodies can be labeled with a variety of detectable labels including,
10 but not limited to, radioisotopes and paramagnetic metals. Therapeutics agents such as small molecule and antibodies or fragments thereof which decrease the concentration and/or activity of a BCSG can also be used in the treatment of diseases characterized by overexpression of BCSG. In these
15 applications, the antibody can be used without or with derivatization to a cytotoxic agent such as a radioisotope, enzyme, toxin, drug or a prodrug.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in
20 the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the
25 disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

DESCRIPTION OF THE INVENTION

The present invention relates to diagnostic assays and
30 methods, both quantitative and qualitative for detecting, diagnosing, monitoring, staging, prognosticating, *in vivo* imaging and treating breast cancer by comparing levels of breast cancer specific genes (BCSGs) with levels of BCSGs in a normal human control. BCSG refers, among other things, to

- 7 -

native proteins expressed by the genes comprising the polynucleotide sequences of BCSG-1 or Gene ID 332369 (SEQ ID NO: 1), BCSG-2 or Gene ID 480489 (SEQ ID NO:2 or 18), BCSG-3 or Gene ID 274731 (SEQ ID NO:3 or 20), BCSG-4 or Gene ID 173388 (SEQ ID NO:4) or BCSG-5 or Clone ID 3040232, Gene ID 411152 (SEQ ID NO:5). Exemplary proteins expressed by genes BCSG-2 and BCSG-3 are depicted herein as SEQ ID NO:19 and SEQ ID NO:21. The genes encoding these proteins (SEQ ID NO:18 and 20) as well as the proteins (SEQ ID NO:19 and 21) have been disclosed in GenBank as Accession No. AF016492.1 (SEQ ID NO:18), AAC27891.1 (SEQ ID NO:19), AF183819 (SEQ ID NO:20) and AAF23614.1 (SEQ ID NO:21). By "BCSG" it is also meant herein variant polynucleotides which, due to degeneracy in genetic coding, comprise variations in nucleotide sequence as compared to SEQ ID NO: 1, 2, 3, 4, 5, 18 or 20 but which still encode the same proteins. The native protein being detected may be whole, a breakdown product, a complex of molecules or chemically modified. In the alternative, what is meant by BCSG as used herein, means the native mRNAs encoded by the genes comprising BCSG-1 or Gene ID 332369 (SEQ ID NO: 1), BCSG-2 or Gene ID 480489 (SEQ ID NO:2 or 18), BCSG-3 or Gene ID 274731 (SEQ ID NO:3 or 20), BCSG-4 or Gene ID 173388 (SEQ ID NO:4) or BCSG-5 or Clone ID 3040232, Gene ID 411152 (SEQ ID NO:5) or it can refer to the actual genes comprising BCSG-1 or Gene ID 332369 (SEQ ID NO: 1), BCSG-2 or Gene ID 480489 (SEQ ID NO:2 or 18), BCSG-3 or Gene ID 274731 (SEQ ID NO:3 or 20), BCSG-4 or Gene ID 173388 (SEQ ID NO:4) or BCSG-5 or Clone ID 3040232, Gene ID 411152 (SEQ ID NO:5) or levels of polynucleotides which are capable of hybridizing under stringent conditions to the antisense sequences of SEQ ID NO: 1, 2, 3, 4, 5, 18 or 20. Such levels are preferably measured in at least one of, cells, tissues and/or bodily fluids, including determination of normal and abnormal levels. Thus, for instance, a diagnostic assay in accordance with the invention for diagnosing over-expression of a BCSG protein

- 8 -

compared to normal control bodily fluids, cells, or tissue samples can be used to diagnose the presence of cancers, and in particular breast cancer. BCSGs may be measured alone in the methods of the invention, or, more preferably, in
5 combination with other diagnostic markers for breast cancer including other BCSGs as described herein. Other breast cancer markers, in addition to BCSGs, useful in the present invention are known to those of skill in the art.

Diagnostic Assays

10 The present invention provides methods for diagnosing the presence of cancer, and in particular breast cancer, by analyzing for changes in levels of BCSG in cells, tissues or bodily fluids from a human patient compared with levels of BCSG in cells, tissues or bodily fluids of preferably the same
15 type from a normal human control, wherein an increase in levels of BCSG in the patient versus the normal human control is associated with the presence of cancer.

Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating
20 the patient being tested has breast cancer is one in which cells, tissues, or bodily fluid levels of a cancer marker, such as BCSG, are at least two times higher, and most preferably are at least five times higher, than in preferably the same cells, tissues, or bodily fluid of a normal human
25 control.

The present invention also provides a method of diagnosing metastatic cancer, and in particular metastatic breast cancer, in a patient having breast cancer which has not yet metastasized. In the method of the present invention, a
30 human cancer patient suspected of having breast cancer which may have metastasized (but which was not previously known to have metastasized) is identified. This is accomplished by a variety of means known to those of skill in the art.

In the present invention, determining the presence of
35 BCSG in cells, tissues, or bodily fluid, is particularly

- 9 -

useful for discriminating between cancers which have not metastasized and cancers which have metastasized. Existing techniques have difficulty discriminating between breast cancer which has metastasized and breast cancer which has not
5 metastasized. However, proper treatment selection is often dependent upon such knowledge.

In the present invention, one of the cancer marker levels measured in cells, tissues, or bodily fluid of a human patient is BCSG. Levels in the human patient are compared with levels
10 of BCSG in preferably the same cells, tissue, or bodily fluid type of a normal human control. That is, if the cancer marker being observed is BCSG in serum, this level is preferably compared with the level of BCSG in serum of a normal human control. An increase in BCSG in the human patient versus the
15 normal human control is associated with a cancer which has metastasized.

Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the cancer in the patient being tested or monitored has
20 metastasized is one in which cells, tissues, or bodily fluid levels of a cancer marker, such as BCSG, are at least two times higher, and more preferably are at least five times higher, than in preferably the same cells, tissues, or bodily fluid of a normal human control.

25 Normal human control as used herein includes a human patient without cancer and/or non cancerous samples from the patient; in the methods for diagnosing metastasis or monitoring for metastasis, normal human control preferably includes samples from a human patient that is determined by
30 reliable methods to have breast cancer which has not metastasized, such as samples from the same patient prior to metastasis.

Staging

The invention also provides a method of staging cancers
35 in a human patient.

- 10 -

The method comprises identifying a human patient having breast cancer and analyzing a sample of cells, tissues, or bodily fluid from such patient for BCSG. The measured BCSG levels are then compared to levels of BCSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein an increase in BCSG levels in the human patient versus the normal human control is associated with breast cancer which is progressing and a decrease in the levels of BCSG is associated with breast cancer which is regressing or in remission.

Monitoring

Further provided is a method of monitoring breast cancer in a human patient for the onset of metastasis. The method comprises identifying a human patient having breast cancer that is not known to have metastasized; periodically analyzing cells, tissues, or bodily fluid from such patient for BCSG; and comparing the BCSG levels in such cells, tissue, or bodily fluid with levels of BCSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein an increase in BCSG levels in the patient versus the normal human control is associated with breast cancer which has metastasized.

Further provided by this invention is a method of monitoring the change in stage of breast cancer. The method comprises identifying a human patient having breast cancer; periodically analyzing cells, tissues, or bodily fluid from such patient for BCSG; and comparing the BCSG levels in such cells, tissue, or bodily fluid with levels of BCSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein an increase in BCSG levels in the patient versus the normal human control is associated with breast cancer which is progressing in stage and a decrease in the levels of BCSG is associated with breast cancer which is regressing in stage or in remission.

- 11 -

Monitoring such patients for onset of metastasis is periodic and preferably done on a quarterly basis. However, this may be performed more or less frequently depending on the cancer, the particular patient, and the stage of the cancer.

5 ***Prognostic Testing and Clinical Trial Monitoring***

The methods described herein can further be utilized as prognostic assays to identify subjects having or at risk of developing a disease or disorder associated with increased levels of BCSG. The present invention provides a method in
10 which a test sample is obtained from a human patient and BCSG is detected. The presence of higher BCSG levels as compared to normal human controls is diagnostic for the human patient being at risk for developing cancer, particularly breast cancer.

15 The effectiveness of therapeutic agents to decrease expression or activity of the BCSGs of the invention can also be monitored by analyzing levels of expression of the BCSGs in a human patient in clinical trials or in *in vitro* screening assays such as in human cells. In this way, the gene
20 expression pattern can serve as a marker, indicative of the physiological response of the human patient, or cells as the case may be, to the agent being tested.

Detection of genetic lesions or mutations

The methods of the present invention can also be used to
25 detect genetic lesions or mutations in BCSG, thereby determining if a human with the genetic lesion is at risk for breast cancer or has breast cancer. Genetic lesions can be detected, for example, by ascertaining the existence of a deletion and/or addition and/or substitution of one or more
30 nucleotides from the BCSGs of this invention, a chromosomal rearrangement of BCSG, aberrant modification of BCSG (such as of the methylation pattern of the genomic DNA), the presence of a non-wild type splicing pattern of a mRNA transcript of BCSG, allelic loss of BCSG, and/or inappropriate post-
35 translational modification of BCSG protein. Methods to detect

- 12 -

such lesions in the BCSG of this invention are known to those of skill in the art.

Assay Techniques

Assay techniques that can be used to determine levels of gene expression, such as BCSG of the present invention, in a sample derived from a human are well-known to those of skill in the art. Such assay methods include radioimmunoassays, reverse transcriptase PCR (RT-PCR) assays, immunohistochemistry assays, *in situ* hybridization assays, competitive-binding assays, Western Blot analyses, ELISA assays and proteomic approaches, two-dimensional gel electrophoresis (2D electrophoresis) and non-gel based approaches such as mass spectrometry or protein interaction profiling. Among these, ELISAs are frequently preferred to diagnose a gene's expressed protein in biological fluids.

An ELISA assay initially comprises preparing an antibody, if not readily available from a commercial source, specific to BCSG, preferably a monoclonal antibody. In addition a reporter antibody generally is prepared which binds specifically to BCSG. The reporter antibody is attached to a detectable reagent such as a radioactive, fluorescent or enzymatic reagent, for example horseradish peroxidase enzyme or alkaline phosphatase.

To carry out the ELISA, antibody specific to BCSG is incubated on a solid support, e.g., a polystyrene dish, that binds the antibody. Any free protein binding sites on the dish are then covered by incubating with a non-specific protein such as bovine serum albumin. Next, the sample to be analyzed is incubated in the dish, during which time BCSG binds to the specific antibody attached to the polystyrene dish. Unbound sample is washed out with buffer. A reporter antibody specifically directed to BCSG and linked to a detectable reagent such as horseradish peroxidase is placed in the dish resulting in binding of the reporter antibody to any monoclonal antibody bound to BCSG. Unattached reporter

- 13 -

antibody is then washed out. Reagents for peroxidase activity, including a colorimetric substrate are then added to the dish. Immobilized peroxidase, linked to BCSG antibodies, produces a colored reaction product. The amount
5 of color developed in a given time period is proportional to the amount of BCSG protein present in the sample. Quantitative results typically are obtained by reference to a standard curve.

A competition assay can also be employed wherein
10 antibodies specific to BCSG are attached to a solid support and labeled BCSG and a sample derived from the patient or human control are passed over the solid support. The amount of label detected which is attached to the solid support can be correlated to a quantity of BCSG in the sample.

15 Using all or a portion of a nucleic acid sequence of a BCSG of the present invention as a hybridization probe, nucleic acid methods can also be used to detect BCSG mRNA as a marker for cancer, and in particular breast cancer. Polymerase chain reaction (PCR) and other nucleic acid
20 methods, such as ligase chain reaction (LCR) and nucleic acid sequence based amplification (NASABA), can be used to detect malignant cells for diagnosis and monitoring of various malignancies. For example, reverse-transcriptase PCR (RT-PCR) is a powerful technique which can be used to detect the
25 presence of a specific mRNA population in a complex mixture of thousands of other mRNA species. In RT-PCR, an mRNA species is first reverse transcribed to complementary DNA (cDNA) with use of the enzyme reverse transcriptase; the cDNA is then amplified as in a standard PCR reaction. RT-PCR can
30 thus reveal by amplification the presence of a single species of mRNA. Accordingly, if the mRNA is highly specific for the cell that produces it, RT-PCR can be used to identify the presence of a specific type of cell.

Hybridization to clones or oligonucleotides arrayed on
35 a solid support (i.e., gridding) can be used to both detect

- 14 -

the expression of and quantitate the level of expression of that gene. In this approach, a cDNA encoding a BCSG gene is fixed to a substrate. The substrate may be of any suitable type including but not limited to glass, nitrocellulose, nylon 5 or plastic. At least a portion of the DNA encoding the BCSG gene is attached to the substrate and then incubated with the analyte, which may be RNA or a complementary DNA (cDNA) copy of the RNA, isolated from the tissue of interest. Hybridization between the substrate bound DNA and the analyte 10 can be detected and quantitated by several means including but not limited to radioactive labeling or fluorescence labeling of the analyte or a secondary molecule designed to detect the hybrid. Quantitation of the level of gene expression can be done by comparison of the intensity of the signal from the 15 analyte compared with that determined from known standards. The standards can be obtained by *in vitro* transcription of the target gene, quantitating the yield, and then using that material to generate a standard curve.

Of the proteomic approaches, 2D electrophoresis is a 20 technique well known to those in the art. Isolation of individual proteins from a sample such as serum is accomplished using sequential separation of proteins by different characteristics usually on polyacrylamide gels. First, proteins are separated by size using an electric 25 current. The current acts uniformly on all proteins, so smaller proteins move farther on the gel than larger proteins. The second dimension applies a current perpendicular to the first and separates proteins not on the basis of size but on the specific electric charge carried by each protein. Since 30 no two proteins with different sequences are identical on the basis of both size and charge, the result of a 2D separation is a square gel in which each protein occupies a unique spot. Analysis of the spots with chemical or antibody probes, or subsequent protein microsequencing can reveal the relative

- 15 -

abundance of a given protein and the identity of the proteins in the sample.

The above tests can be carried out on samples derived from a variety cells, bodily fluids and/or tissue extracts
5 (homogenates or solubilized tissue) obtained from the patient including tissue biopsy and autopsy material. Bodily fluids useful in the present invention include blood, urine, saliva, or any other bodily secretion or derivative thereof. Blood can include whole blood, plasma, serum, or any derivative of
10 blood.

In Vivo Targeting of BCSGs/Breast Cancer Therapy

Identification of BCSGs is also useful in the rational design of new therapeutics for imaging and treating cancers, and in particular breast cancer. For example, in one
15 embodiment, antibodies which specifically bind to BCSGs can be raised and used *in vivo* in patients' suspected of suffering from cancer. Antibodies which specifically bind a BCSG can be injected into a patient suspected of having cancer for diagnostic and/or therapeutic purposes. The preparation and
20 use of antibodies for *in vivo* diagnosis is well known in the art. For example, antibody-chelators labeled with Indium-111 have been described for use in the radioimmunoscintigraphic imaging of carcinoembryonic antigen expressing tumors (Sumerdon et al. Nucl. Med. Biol. 1990 17:247-254). In
25 particular, these antibody-chelators have been used in detecting tumors in patients suspected of having recurrent colorectal cancer (Griffin et al. J. Clin. Onc. 1991 9:631-640). Antibodies with paramagnetic ions as labels for use in magnetic resonance imaging have also been described (Lauffer,
30 R.B. Magnetic Resonance in Medicine 1991 22:339-342). Antibodies directed against BCSGs can be used in a similar manner. Labeled antibodies which specifically bind a BCSG can be injected into patients suspected of having breast cancer for the purpose of diagnosing or staging of the disease status
35 of the patient. The label used will be selected in accordance

- 16 -

with the imaging modality to be used. For example, radioactive labels such as Indium-111, Technetium-99m or Iodine-131 can be used for planar scans or single photon emission computed tomography (SPECT). Positron emitting
5 labels such as Fluorine-19 can be used in positron emission tomography. Paramagnetic ions such as Gadolinium (III) or Manganese (II) can be used in magnetic resonance imaging (MRI). Localization of the label permits determination of the spread of the cancer. The amount of label within an organ or
10 tissue also allows determination of the presence or absence of cancer in that organ or tissue.

For patients diagnosed with cancer, and in particular breast cancer, injection of an antibody which specifically binds a BCSG can also have a therapeutic benefit. The
15 antibody may exert its therapeutic effect alone. Alternatively, the antibody can be conjugated to a cytotoxic agent such as a drug, toxin or radionuclide to enhance its therapeutic effect. Drug monoclonal antibodies have been described in the art for example by Garnett and Baldwin,
20 Cancer Research 1986 46:2407-2412. The use of toxins conjugated to monoclonal antibodies for the therapy of various cancers has also been described by Pastan et al. Cell 1986 47:641-648. Yttrium-90 labeled monoclonal antibodies have been described for maximization of dose delivered to the tumor
25 while limiting toxicity to normal tissues (Goodwin and Meares Cancer Supplement 1997 80:2675-2680). Other cytotoxic radionuclides including, but not limited to Copper-67, Iodine-131 and Rhenium-186 can also be used for labeling of antibodies against BCSG.

30 Antibodies which can be used in these *in vivo* methods include polyclonal, monoclonal and omniclonal antibodies and antibodies prepared via molecular biology techniques. Antibody fragments and aptamers and single-stranded oligonucleotides such as those derived from an *in vitro*

- 17 -

evolution protocol referred to as SELEX and well known to those skilled in the art can also be used.

Screening Assays

The present invention also provides methods for
5 identifying modulators which bind to BCSG protein or have a modulatory effect on the expression or activity of BCSG protein. Modulators which decrease the expression or activity of BCSG protein are believed to be useful in treating breast cancer. Such screening assays are known to those of skill in
10 the art and include, without limitation, cell-based assays and cell free assays.

Small molecules predicted via computer imaging to specifically bind to regions of BCSG can also be designed, synthesized and tested for use in the imaging and treatment
15 of breast cancer. Further, libraries of molecules can be screened for potential anticancer agents by assessing the ability of the molecule to bind to the BCSGs identified herein. Molecules identified in the library as being capable of binding to BCSG are key candidates for further evaluation
20 for use in the treatment of breast cancer. In a preferred embodiment, these molecules will downregulate expression and/or activity of BCSG in cells.

Adoptive Immunotherapy and Vaccines

Adoptive immunotherapy of cancer refers to a therapeutic
25 approach in which immune cells with an antitumor reactivity are administered to a tumor-bearing host, with the aim that the cells mediate either directly or indirectly, the regression of an established tumor. Transfusion of lymphocytes, particularly T lymphocytes, falls into this
30 category and investigators at the National Cancer Institute (NCI) have used autologous reinfusion of peripheral blood lymphocytes or tumor-infiltrating lymphocytes (TIL), T cell cultures from biopsies of subcutaneous lymph nodules, to treat several human cancers (Rosenberg, S. A., U.S. Patent No.

- 18 -

4,690,914, issued Sep. 1, 1987; Rosenberg, S. A., et al., 1988, N. England J. Med. 319:1676-1680).

The present invention relates to compositions and methods of adoptive immunotherapy for the prevention and/or treatment
5 of primary and metastatic breast cancer in humans using macrophages sensitized to the antigenic BCSG molecules, with or without non-covalent complexes of heat shock protein (hsp). Antigenicity or immunogenicity of the BCSG is readily confirmed by the ability of the BCSG protein or a fragment
10 thereof to raise antibodies or educate naive effector cells, which in turn lyse target cells expressing the antigen (or epitope).

Cancer cells are, by definition, abnormal and contain proteins which should be recognized by the immune system as
15 foreign since they are not present in normal tissues. However, the immune system often seems to ignore this abnormality and fails to attack tumors. The foreign BCSG proteins that are produced by the cancer cells can be used to reveal their presence. The BCSG is broken into short fragments, called
20 tumor antigens, which are displayed on the surface of the cell. These tumor antigens are held or presented on the cell surface by molecules called MHC, of which there are two types: class I and II. Tumor antigens in association with MHC class I molecules are recognized by cytotoxic T cells while antigen-
25 MHC class II complexes are recognized by a second subset of T cells called helper cells. These cells secrete cytokines which slow or stop tumor growth and help another type of white blood cell, B cells, to make antibodies against the tumor cells.

30 In adoptive immunotherapy, T cells or other antigen presenting cells (APCs) are stimulated outside the body (ex vivo), using the tumor specific BCSG antigen. The stimulated cells are then reinfused into the patient where they attack the cancerous cells. Research has shown that using both
35 cytotoxic and helper T cells is far more effective than using

- 19 -

either subset alone. Additionally, the BCSG antigen may be complexed with heat shock proteins to stimulate the APCs as described in U.S. Patent No. 5,985,270.

The APCs can be selected from among those antigen
5 presenting cells known in the art including, but not limited to, macrophages, dendritic cells, B lymphocytes, and a combination thereof, and are preferably macrophages. In a preferred use, wherein cells are autologous to the individual, autologous immune cells such as lymphocytes, macrophages or
10 other APCs are used to circumvent the issue of whom to select as the donor of the immune cells for adoptive transfer. Another problem circumvented by use of autologous immune cells is graft versus host disease which can be fatal if unsuccessfully treated.

15 In adoptive immunotherapy with gene therapy, DNA of the BCSG can be introduced into effector cells similarly as in conventional gene therapy. This can enhance the cytotoxicity of the effector cells to tumor cells as they have been manipulated to produce the antigenic protein resulting in
20 improvement of the adoptive immunotherapy.

BCSG antigens of this invention are also useful as components of breast cancer vaccines. The vaccine comprises an immunogenically stimulatory amount of an BCSG antigen. Immunogenically stimulatory amount refers to that amount of
25 antigen that is able to invoke the desired immune response in the recipient for the amelioration, or treatment of breast cancer. Effective amounts may be determined empirically by standard procedures well known to those skilled in the art.

The BCSG antigen may be provided in any one of a number
30 of vaccine formulations which are designed to induce the desired type of immune response, e.g., antibody and/or cell mediated. Such formulations are known in the art and include, but are not limited to, formulations such as those described in U.S. Patent 5,585,103. Vaccine formulations of the present

- 20 -

invention used to stimulate immune responses can also include pharmaceutically acceptable adjuvants.

EXAMPLES

The present invention is further described by the following examples. These examples are provided solely to illustrate the invention by reference to specific embodiments. These exemplifications, while illustrating certain specific aspects of the invention, do not portray the limitations or circumscribe the scope of the disclosed invention.

10 Example 1: Identification of BCSGs via CLASP

Identification of BCSGs (Breast Cancer Specific Genes) was carried out by a systematic analysis of data in the LIFESEQ Gold (LSGold) database available from Incyte Pharmaceuticals, Palo Alto, CA using the data mining Cancer Leads Automatic Search Package (CLASP) developed by diaDexus LLC, Santa Clara CA.

The CLASP performs the following steps:

(1) Selection of highly expressed organ specific genes based on the abundance level of the corresponding EST in the targeted organ versus all the other organs.

(2) Analysis of the expression level of each highly expressed organ specific gene in normal, tumor tissue, disease tissue and tissue libraries associated with tumor or disease.

(3) Selection of the candidates wherein component ESTs are exclusively or more frequently found in tumor libraries.

The CLASP allows the identification of highly expressed organ and cancer specific genes. A final manual in depth evaluation is then performed to finalize the Organ Cancer Specific Genes (OCSGs) selection. Table 1 provides the BCSGs of the present invention identified using CLASP.

- 21 -

Table 1: BCSGs

BCSG	SEQ ID NO:	LSGold Clone ID	LSGold Gene ID
BCSG-1	1	none	332369
BCSG-2	2 or 18	none	480489
5 BCSG-3	3 or 20	none	274731
BCSG-4	4	none	173388
BCSG-5	5	3040232	411152

Example 2: Determination of mRNA expression of BCSG-5

The mRNA expression level of BCSG, BCSG-5 (SEQ ID NO:5, Clone ID 3040232, Gene ID 411152), also referred to as MAM009, in different tissues was analyzed using Real-Time quantitative PCR. The results presented here for BCSG-5 support the usage of CLASP as a tool for identifying cancer diagnostic markers.

These experiments were carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. Routine molecular biology techniques were carried out as described in standard laboratory manuals, such as Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989).

Real-Time quantitative PCR with fluorescent Taqman probes is a quantitation detection system utilizing the 5'-3' nuclease activity of Taq DNA polymerase. The method uses an internal fluorescent oligonucleotide probe (Taqman) labeled with a 5' reporter dye and a downstream, 3' quencher dye. During PCR, the 5'-3' nuclease activity of Taq DNA polymerase releases the reporter, whose fluorescence can then be detected by the laser detector of the Model 7700 Sequence Detection System (PE Applied Biosystems, Foster City, CA, USA).

Amplification of an endogenous control is used to standardize the amount of sample RNA added to the reaction and normalize for Reverse Transcriptase (RT) efficiency. Either

- 22 -

cyclophilin, glyceraldehyde-3-phosphate dehydrogenase (GAPDH) or 18S ribosomal RNA (rRNA) was used as this endogenous control. To calculate relative quantitation between all the samples studied, the target RNA levels for one sample were
5 used as the basis for comparative results (calibrator). Quantitation relative to the "calibrator" can be obtained using the standard curve method or the comparative method (User Bulletin #2: ABI PRISM 7700 Sequence Detection System).

The tissue distribution, and the level of the target gene
10 were determined for every sample in normal and cancer tissue. Total RNA was extracted from normal tissues, cancer tissues, and from cancers and the corresponding matched adjacent tissues. Subsequently, first strand cDNA was prepared with reverse transcriptase and the polymerase chain reaction was
15 done using primers and Taqman probe specific to each target gene. The results were analyzed using the ABI PRISM 7700 Sequence Detector. The absolute numbers are relative levels of expression of the target gene in a particular tissue compared to the calibrator tissue.

20 Primers used for expression analysis include:

BCSG-5 forward:

ACCCCATTTAGCCTGCCAT (SEQ ID NO:6)

BCSG-5 reverse:

ATGGGAGTATCTCATCTGCTCTCA (SEQ ID NO:7)

25 Q-PCR probe:

TGTTTGTTTCATTCTTCAATTCCAAGGCTTT (SEQ ID NO:8)

The absolute numbers depicted in Table 2 are relative levels of expression of BCSG-5 in 12 normal different tissues. All the values are compared to normal testis (calibrator).
30 These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

- 23 -

Table 2: Relative Levels of BCSG-5 Expression in Pooled Samples

	Tissue	NORMAL
5	Brain	0.00
	Heart	0.00
	Kidney	0.00
	Liver	0.00
	Lung	0.00
	Mammary gland	106.15
10	Muscle	0.00
	Prostate	0.00
	Small Intestine	0.00
	Testis	1.00
	Thymus	0.00
15	Uterus	0.00

The relative levels of expression in Table 2 show that BCSG-5 mRNA expression is detected in the pool of normal mammary gland and in testis but not in the other 10 normal tissue pools analyzed. The level of expression in mammary gland is more than 100 fold higher than in testis. These results demonstrate that BCSG-5 mRNA expression is highly specific for mammary gland tissue and is also found in testis. Expression in a male specific tissue is not relevant in detecting cancer in female specific tissues.

The absolute numbers in Table 2 were obtained analyzing pools of samples of a particular tissue from different individuals. They can not be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 3.

The absolute numbers depicted in Table 3 are relative levels of expression of BCSG-5 in 78 pairs of matching samples. All the values are compared to normal testis (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual. In addition, 2 unmatched cancer samples (from ovary) and 2 unmatched normal samples (from ovary) were also tested.

- 24 -

Table 3: Relative Levels of BCSG-5 Expression in Individual Samples

	Sample ID	Tissue	Cancer	Matching Normal Adjacent	Normal
5	MamS621	Mammary Gland 1	60.37	0.00	
	MamS516	Mammary Gland 2	1.97	1.09	
	MamS079	Mammary Gland 3	2.31	2.74	
	Mam517	Mammary Gland 4	3.42	2.71	
	Mam59X	Mammary Gland 5	0.47	9.56	
10	MamS127	Mammary Gland 6	0.00	2.22	
	MamB011X	Mammary Gland 7	2.52	25.28	
	Mam522	Mammary Gland 8	109.66	2.67	
	Mam51DN	Mammary Gland 9	11.71	169.77	
	Mam19DN	Mammary Gland 10	369.64	28.24	
15	MamS123	Mammary Gland 11	0.10	1.21	
	MamS997	Mammary Gland 12	8.80	2.29	
	Mam162X	Mammary Gland 13	7.67	1.08	
	Mam220	Mammary Gland 14	11.50	53.60	
	Mam699F	Mammary Gland 15	0.52	3.48	
20	Mam42DN	Mammary Gland 16	1.39	3.54	
	Mam76DN	Mammary Gland 17	300.03	84.71	
	MamS854	Mammary Gland 18	2.77	2.64	
	MamS967	Mammary Gland 19	892.68	4.46	
	Mam986	Mammary Gland 20	14.40	19.27	
25	MamS699	Mammary Gland 21	2.24	1.43	
	Mam355	Mammary Gland 22	223.37	0.00	
	MamA06X	Mammary Gland 23	1220.50	2.26	
	MamS570	Mammary Gland 24	0.00	120.39	
	MamS918	Mammary Gland 25	181.43	60.30	
30	End12XA	Endometrium 1	0.00	0.00	

- 25 -

	End28XA	Endometrium 2	0.00	0.00	
	End3AX	Endometrium 3	0.00	0.00	
	End4XA	Endometrium 4	0.00	0.00	
	End5XA	Endometrium 5	0.00	0.00	
5	End10479	Endometrium 6	0.00	0.00	
	End65RA	Endometrium 7	0.00	0.00	
	End68X	Endometrium 8	0.00	0.00	
	CvxNKS18	Cervix 1	0.00	0.00	
	CvxNKS54	Cervix 2	0.00	0.00	
10	CvxNK23	Cervix 3	0.00	0.00	
	CvxNK24	Cervix 4	0.00	0.00	
	CvxKS52	Cervix 5	0.00	0.00	
	CvxKS83	Cervix 6	0.00	0.00	
	Utr141XO	Uterus 1	0.00	0.00	
15	Utr135XO	Uterus 2	0.00	0.00	
	Utr23XU	Uterus 3	0.00	0.00	
	Utr85XU	Uterus 4	0.00	0.00	
	LngC20X	Lung 1	0.00	0.00	
	LngSQ56	Lung 2	0.00	0.00	
20	Lng90X	Lung 3	0.00	0.00	
	LngAC11	Lung 4	0.00	0.00	
	Pro101XB	Prostate 1	0.00	0.00	
	Pro23B	Prostate 2	0.00	0.00	
	Skn448S	Skin 1	0.00	0.00	
25	Skn784S	Skin 2	0.00	0.00	
	ClnSG45	Colon 1	0.00	0.00	
	ClnTX01	Colon 2	0.00	0.00	
	ClnAS46	Colon 3	0.00	0.00	
	ClnAS67	Colon 4	0.00	0.00	
30	BldTR17	Bladder 1	0.00	0.00	

- 26 -

	Bld66X	Bladder 2	0.00	0.00	
	Kid11XD	Kidney 1	0.00	0.00	
	Kid5XD	Kidney 2	0.00	0.00	
	Kid109XD	Kidney 3	0.00	0.00	
5	Liv532L	Liver 1	0.00	0.00	
	Liv175L	Liver 2	0.00	0.00	
	Liv187L	Liver 3	0.00	0.00	
	OvrG010	Ovary 1	0.00	0.00	
	Ovr10050	Ovary 2	0.00		
10	Ovr1028	Ovary 3	0.00		
	Ovr103X	Ovary 4	0.00	0.00	
	Ovr18GA	Ovary 5			0.00
	Ovr206I	Ovary 6			0.00
	Pan92X	Pancreas 1	0.00	0.00	
15	PanC044	Pancreas 2	0.00	0.00	
	SmIH89	Small Intestine 1	0.00	0.00	
	SmI21XA	Small Intestine 2	0.00	0.00	
	Sto15S	Stomach 1	0.00	0.00	
	StoAC44	Stomach 2	0.00	0.00	
20	Sto288S	Stomach 3	0.00	0.00	
	Sto531S	Stomach 4	0.00	0.00	
	Thr644T	Thyroid 1	0.00	0.00	
	Thr145T	Thyroid 2	0.00	0.00	
	Thr939T	Thyroid 3	0.00	0.00	
25	Tst39X	Testis 1	0.00	0.00	
	Tst663T	Testis 2	0.00	0.00	

0.00= Negative

Among 160 samples in Table 3 representing 17 different tissues significant expression is seen only in mammary gland
 30 tissues. These results confirm the tissue specificity results obtained with normal samples shown in Table 2. Table 2 and

- 27 -

Table 3 represent a combined total of 172 samples in 21 human tissue types. One hundred and twenty samples representing 20 different tissue types excluding mammary gland had no detectable level of BCSG-5 mRNA. Other than mammary gland, 5 BCSG-5 is detected only in one tissue type (testis) and then only in the pooled tissue sample (Table 2) but not in the matched testis cancer samples (Table 3; testis 1 and 2).

Comparisons of the level of mRNA expression in breast cancer samples and the normal adjacent tissue from the same 10 individuals are shown in Table 3. BCSG-5 is expressed at higher levels in 12 of 25 (48%) cancer samples (mammary gland 1, 2, 8, 10, 12, 13, 17, 19, 21, 22, 23 and 25) compared to normal adjacent tissue.

Altogether, the high level of tissue specificity, plus 15 the mRNA overexpression in 48% of the mammary gland matching samples tested are indicative of BCSG-5, and more generally BCSGs selected by CLASP, being good diagnostic markers for breast cancer.

Example 3: Determination of mRNA expression of BCSG-1
20 The mRNA expression level of BCSG, BCSG-1 (SEQ ID NO:1, Gene ID 332369), also referred to as MAM014 were also determined in accordance with methods as set forth in Example 2.

Real-Time quantitative PCR was done using the following 25 primers:

BCSG-1 forward:

5' GCCCATTAGCACCCAGATAAT 3' (SEQ ID NO:9)

BCSG-2 reverse:

5' GCCAACCCTTCACCTAAGAAA 3' (SEQ ID NO:10)

30 Q-PCR probe

5' CTCCCCACTGTACAAAGATTTTCCAGGATG 3' (SEQ ID NO:11)

The absolute numbers depicted in Table 4 are relative levels of expression of BCSG-1 in 37 normal samples from 25 different tissues. All the values are compared to normal

- 28 -

kidney (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals; except for the blood samples that they are normal samples from a single individual.

5 Table 4: Relative Levels of BCSG-1 Expression in Pooled Samples

	Tissue	NORMAL
	Adrenal Gland	1.09
	Bladder	0.05
10	Brain	24.00
	Cervix	3.84
	Colon	0.00
	Endometrium	10.41
	Esophagus	0.18
15	Heart	0.01
	Kidney	1.00
	Liver	0.02
	Lung	4.35
	Mammary	1.19
20	Muscle	0.09
	Ovary	23.51
	Pancreas	0.86
	Prostate	7.75
	Rectum	0.33
25	Small Intestine	0.85
	Spleen	17.51
	Stomach	2.42
	Testis	111.04
	Thymus	9.95
30	Trachea	6.43
	Uterus	0.68
	Blood 1	34.42
	Blood 2	0.00
	Blood 3	21.19
35	Blood 4	25.19
	Blood 5	51.09
	Blood 6	1144.10
	Blood 7	59.10
	Blood 8	60.13
40	Blood 9	37.53
	Blood 10	0.00
	Blood 11	15.30
	Blood 12	0.00
	Blood 13	0.00

45 The relative levels of expression in Table 4 show that BCSG-1 mRNA expression is detected in the pool of normal mammary gland as well as in the other normal tissue analyzed.

- 29 -

The absolute numbers in Table 5 were obtained analyzing pools of samples of a particular tissue from different individuals, except for the blood samples. They can not be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 5.

The absolute numbers depicted in Table 5 are relative levels of expression of BCSG-1 in 77 pairs of matching samples. All the values are compared to normal kidney (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual. In addition, 3 unmatched cancer samples (from ovary) and 3 unmatched normal samples (from ovary) were also tested.

Table 5: Relative Levels of BCSG-1 Expression in Individual Samples

Sample ID	Tissue	Cancer	Matching Normal Adjacent	Normal
MamS621	Mammary Gland 1	7.48	0.00	
MamS516	Mammary Gland 2	0.90	0.00	
Mam173M	Mammary Gland 3	3.96	0.00	
Mam497M	Mammary Gland 4	9.71	0.00	
MamS079	Mammary Gland 5	1.72	0.00	
Mam517	Mammary Gland 6	8.88	3.35	
Mam726M	Mammary Gland 7	3.04	0.00	
Mam59X	Mammary Gland 8	7.01	15.73	
MamS127	Mammary Gland 9	24.59	0.00	
MamB011X	Mammary Gland 10	8.43	1.14	
MamS22	Mammary Gland 11	14.55	0.00	
Mam15DN	Mammary Gland 12	4.16	0.00	
Mam51DN	Mammary Gland 13	32.90	3.11	
Mam976M	Mammary Gland 14	6.17	0.00	
Mam543M	Mammary Gland 15	34.42	0.32	

- 30 -

	Mam245M	Mammary Gland 16	10.82	0.00	
	MamS123	Mammary Gland 17	0.37	0.00	
	MamS997	Mammary Gland 18	0.56	0.00	
	Mam162X	Mammary Gland 19	6.09	0.52	
5	Mam220	Mammary Gland 20	2.08	0.58	
	Mam699F	Mammary Gland 21	6.75	6.36	
	Mam42DN	Mammary Gland 22	10.16	0.00	
	Mam76DN	Mammary Gland 23	31.23	4.68	
	MamS854	Mammary Gland 24	6.11	0.00	
10	MamS967	Mammary Gland 25	86.22	0.00	
	Mam986	Mammary Gland 26	13.36	9.00	
	MamS699	Mammary Gland 27	4.52	0.00	
	Mam355	Mammary Gland 28	107.38	0.00	
	MamA06X	Mammary Gland 29	43.26	0.00	
15	MamS570	Mammary Gland 30	68.36	64.22	
	MamS918	Mammary Gland 31	2.49	0.86	
	Bld66X	Bladder	0.00	3.24	
	ClnTX01	Colon 1	0.55	0.00	
	ClnAS43	Colon 2	1.11	0.00	
20	ClnAS49	Colon 3	0.69	0.60	
	ClnRS45	Colon 4	0.00	0.00	
	CvxNK24	Cervix 1	2.53	0.69	
	CvxNKS54	Cervix 2	1.71	0.54	
	CvxNK23	Cervix 3	0.34	0.00	
25	CvxNKS81	Cervix 4	0.00	0.00	
	End5XA	Endometrium 1	1.16	2.85	
	End8911	Endometrium 2	2.62	1.65	
	End8963	Endometrium 3	6.50	0.00	
	End28XA	Endometrium 4	1.75	1.33	
30	End65RA	Endometrium 5	0.45	0.00	

- 31 -

	End12XA	Endometrium 6	34.80	3.55	
	End3AX	Endometrium 7	0.00	0.00	
	Kid11XD	Kidney 1	0.94	1.42	
	Kid124D	Kidney 2	1.35	0.00	
5	Liv532L	Liver 1	0.33	0.00	
	Liv390L	Liver 2	0.66	0.00	
	LngSQ56	Lung 1	0.00	0.00	
	Lng223L	Lung 2	0.20	0.00	
	LngLC71	Lung 3	7.67	8.20	
10	LngAC90	Lung 4	11.70	0.98	
	Lng75XC	Lung 5	0.00	0.00	
	OvrA082	Ovary 1	21.33	42.96	
	OvrA082	Ovary 2	52.68	186.62	
	Ovr103X	Ovary 3	44.88	17.67	
15	Ovr10050	Ovary 4	6.89		
	Ovr1028	Ovary 5	0.44		
	Ovr10400	Ovary 6	0.80		
	Ovr18GA	Ovary 7			6.63
	Ovr206I	Ovary 8			1.46
20	Ovr20GA	Ovary 9			5.96
	Pan92X	Pancreas 1	0.00	0.00	
	Pan77X	Pancreas 2	0.24	0.00	
	Pro23B	Prostate 1	0.80	0.00	
	Pro13XB	Prostate 2	0.05	17.75	
25	Skn448S	Skin 1	0.00	0.00	
	Skn784S	Skin 2	0.13	0.11	
	SmIntH89	Small Intestine 1	0.00	0.00	
	Sto264S	Stomach 1	0.80	1.15	
	Sto15S	Stomach 2	0.23	2.29	
30	Sto27S	Stomach 3	1.07	1.35	

- 32 -

	Thr644T	Thyroid 1	0.00	0.00	
	Thr143T	Thyroid 2	0.58	0.00	
	Tst663T	Testis 1	5.46	1.80	
	Tst647T	Testis 2	1.27	5.68	
5	Utr23XU	Uterus 1	9.38	1.17	
	Utr85XU	Uterus 2	4.36	2.17	
	Utr141XO	Uterus 3	0.00	0.00	
	Utr135XO	Uterus 4	7.53	10.10	

0.00= Negative

10 Table 5 represents 160 samples in 17 different tissues. Table 4 and Table 5 represent a combined total of 197 samples in 27 human tissue types. Comparisons of the level of mRNA expression in breast cancer samples and the normal adjacent tissue from the same individuals are shown

15 in Table 5. BCSG-1 is expressed at higher levels in 27 of 30 (90%) cancer samples (mammary gland 1-7, 9-20, 22-25, 27-29, and 31) compared to normal adjacent tissue.

Example 4: Determination of mRNA expression of BCSG-2

The mRNA expression level of BCSG, BCSG-2 (SEQ ID NO:2 or 18, Gene ID 480489), also referred to as MAM013 were

20 also determined in accordance with methods as set forth in Example 2.

Real-Time quantitative PCR was done using the following primers:

25 BCSG-2 forward:

5' CCTGGAGTTTCAATTCCTCA 3' (SEQ ID NO:12)

BCSG-2 reverse:

5' CCCCAGAGAAAACACCACAA 3' (SEQ ID NO:13)

Q-PCR probe

30 5' ACTCCTCCATTTCCTTAGGTAGGGTTTG 3' (SEQ ID NO:14)

The absolute numbers depicted in Table 6 are relative levels of expression of BCSG-2 in 37 normal samples from 25

- 33 -

different tissues. All the values are compared to normal liver (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals, except for the blood samples that they are normal samples from a single individual.

Table 6: Relative Levels of BCSG-2 Expression in Pooled Samples

	Tissue	NORMAL
10	Adrenal Gland	0.00
	Bladder	0.00
	Brain	0.00
	Cervix	0.00
	Colon	0.00
15	Endometrium	0.33
	Esophagus	0.00
	Heart	0.02
	Kidney	0.28
	Liver	1.00
20	Lung	0.07
	Mammary	20.39
	Muscle	0.00
	Ovary	0.00
	Pancreas	0.05
25	Prostate	0.26
	Rectum	0.00
	Small Intestine	0.00
	Spleen	0.00
	Stomach	0.12
30	Testis	1.55
	Thymus	0.55
	Trachea	1.23
	Uterus	0.00
35	Blood 1	0.00
	Blood 2	0.00
	Blood 3	0.00
	Blood 4	0.00
	Blood 5	6.17
	Blood 6	0.00
40	Blood 7	0.00
	Blood 8	0.00
	Blood 9	16.97
	Blood 10	0.00
	Blood 11	77.98
45	Blood 12	0.00
	Blood 13	0.00

- 34 -

The relative levels of expression in Table 6 show that BCSG-2 mRNA expression is detected in the pool of normal mammary gland. The level of expression is higher than in the other tissues with the exception of two blood samples.

5 The absolute numbers in Table 6 were obtained analyzing pools of samples of a particular tissue from different individuals, except for the blood samples. They can not be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in
10 Table 7.

The absolute numbers depicted in Table 7 are relative levels of expression of BCSG-2 in 76 pairs of matching samples. All the values are compared to normal liver (calibrator). A matching pair is formed by mRNA from the
15 cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual. In addition, 3 unmatched cancer samples (from ovary) and 3 unmatched normal samples (from ovary) were also tested.

20 **Table 7: Relative Levels of BCSG-2 Expression in Individual Samples**

Sample ID	Tissue	Cancer	Matching Normal Adjacent	Normal
MamS621	Mammary Gland 1	0.00	0.00	
25 Mam173M	Mammary Gland 2	7.21	0.00	
Mam497M	Mammary Gland 3	1634.92	31.12	
MamS079	Mammary Gland 4	42.56	1.39	
Mam517	Mammary Gland 5	2.42	4.21	
Mam726M	Mammary Gland 6	0.00	6.75	
30 Mam59X	Mammary Gland 7	0.00	16.11	
MamS127	Mammary Gland 8	5.31	0.00	
MamB011X	Mammary Gland 9	0.87	182.28	

- 35 -

	Mam522	Mammary Gland 10	480.47	0.18	
	Mam19DN	Mammary Gland 11	4.79	0.22	
	Mam51DN	Mammary Gland 12	19.49	72.76	
	Mam976M	Mammary Gland 13	62.25	0.00	
5	Mam543M	Mammary Gland 14	103.97	0.00	
	Mam245M	Mammary Gland 15	49.01	615.24	
	MamS123	Mammary Gland 16	0.42	1.27	
	MamS997	Mammary Gland 17	0.24	0.66	
	Mam162X	Mammary Gland 18	0.45	1.39	
10	Mam220	Mammary Gland 19	0.00	0.00	
	Mam699F	Mammary Gland 20	0.00	12.38	
	Mam42DN	Mammary Gland 21	44.48	11.47	
	Mam76DN	Mammary Gland 22	9.32	26.26	
	MamS854	Mammary Gland 23	6.50	103.61	
15	MamS967	Mammary Gland 24	3.36	5.13	
	Mam986	Mammary Gland 25	7.67	65.12	
	MamS699	Mammary Gland 26	1.68	11.63	
	Mam355	Mammary Gland 27	1.32	0.00	
	MamA06X	Mammary Gland 28	1.73	0.26	
20	MamS570	Mammary Gland 29	0.00	194.69	
	MamS918	Mammary Gland 30	0.07	0.13	
	Bld66X	Bladder	0.00	0.00	
	ClnTX01	Colon 1	0.00	0.00	
	ClnAS43	Colon 2	0.00	0.00	
25	ClnAS49	Colon 3	0.00	0.00	
	ClnRS45	Colon 4	0.00	0.01	
	CvxNK24	Cervix 1	0.00	0.00	
	CvxNKS54	Cervix 2	0.00	0.00	
	CvxNK23	Cervix 3	0.02	0.00	
30	CvxNKS81	Cervix 4	0.00	0.00	

- 36 -

	End5XA	Endometrium 1	0.00	0.00	
	End8911	Endometrium 2	0.00	0.00	
	End8963	Endometrium 3	0.00	0.00	
	End28XA	Endometrium 4	0.05	0.00	
5	End65RA	Endometrium 5	0.00	0.00	
	End12XA	Endometrium 6	0.23	0.00	
	End3AX	Endometrium 7	0.00	0.05	
	Kid11XD	Kidney 1	0.04	0.00	
	Kid124D	Kidney 2	0.37	0.00	
10	Liv532L	Liver 1	0.00	2.02	
	Liv390L	Liver 2	0.08	0.56	
	Lng223L	Lung 1	0.00	0.00	
	LngLC71	Lung 2	0.00	0.00	
	LngSQ56	Lung 3	0.00	0.00	
15	LngAC90	Lung 4	0.00	0.00	
	Lng75XC	Lung 5	0.00	0.00	
	OvrA082	Ovary 1	0.00	0.00	
	OvrA082	Ovary 2	0.00	0.00	
	Ovr103X	Ovary 3	0.07	0.00	
20	Ovr10050	Ovary 4	1.09		
	Ovr1028	Ovary 5	0.00		
	Ovr10400	Ovary 6	0.00		
	Ovr18GA	Ovary 7			0.00
	Ovr206I	Ovary 8			0.00
25	Ovr20GA	Ovary 9			0.00
	Pan92X	Pancreas 1	0.00	0.00	
	Pan77X	Pancreas 2	1.91	1.11	
	Pro23B	Prostate 1	0.01	0.00	
	Pro13XB	Prostate 2	0.00	0.00	
30	Skn448S	Skin 1	0.00	0.00	

- 37 -

	Skn784S	Skin 2	0.00	0.00	
	SmIntH89	Small Intestine 1	0.00	0.02	
	Sto264S	Stomach 1	0.00	0.00	
	Sto15S	Stomach 2	0.04	0.00	
5	Sto27S	Stomach 3	0.00	0.00	
	Thr644T	Thyroid 1	0.00	0.12	
	Thr143T	Thyroid 2	0.03	3.73	
	Tst663T	Testis 1	0.09	0.00	
	Tst647T	Testis 2	0.55	0.00	
10	Utr23XU	Uterus 1	0.00	0.00	
	Utr85XU	Uterus 2	0.00	2.85	
	Utr141XO	Uterus 3	0.04	0.00	
	Utr135XO	Uterus 4	0.17	0.12	

0.00= Negative

15 Table 7 represents 158 samples in 17 different tissues. Table 6 and Table 7 represent a combined total of 195 samples in 25 human tissue types. Comparisons of the level of mRNA expression in breast cancer samples and the normal adjacent tissue from the same individuals are shown

20 in Table 7. BCSG-2 is expressed at higher levels in 11 of 30 (37%) cancer samples (mammary gland 2-4, 8, 10, 11, 13, 14, 21, 27, 28) compared to normal adjacent tissue.

Example 5: Determination of mRNA expression of BCSG-3

The mRNA expression level of BCSG, BCSG-3 (SEQ ID NO:3 or 20, Gene ID 274731), also referred to as MAM017 were also determined in accordance with methods as set forth in Example 2.

Real-Time quantitative PCR was done using the following primers:

30 BCSG-3 forward:

5' GAGCACTTCCTTTTGGTTTTTC 3' (SEQ ID NO:15)

BCSG-3 reverse:

- 38 -

5' GCCCTAGCATATTCCAGAAGTTC 3' (SEQ ID NO:16)

Q-PCR probe

5' TAGACAGTGGGCTCACATGTTCTGATAGTG 3' (SEQ ID NO:17)

The absolute numbers depicted in Table 8 are relative levels of expression of BCSG-3 in 36 normal samples from 25 different tissues. All the values are compared to normal prostate (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals, except for 10 the blood samples that they are normal samples from a single individual.

Table 8: Relative Levels of BCSG-3 Expression in Pooled Samples

	Tissue	NORMAL
15	Adrenal Gland	0.16
	Bladder	0.02
	Brain	0.12
	Cervix	1.41
	Colon	0.01
20	Endometrium	3.77
	Esophagus	0.03
	Heart	0.02
	Kidney	0.07
	Liver	0.00
25	Lung	0.59
	Mammary	7.67
	Muscle	0.08
	Ovary	0.94
	Pancreas	0.14
30	Prostate	1.00
	Rectum	0.13
	Small Intestine	0.05
	Spleen	0.89
	Stomach	0.17
35	Testis	0.20
	Thymus	0.56
	Trachea	0.39
	Uterus	1.22
	Blood 1	1.91
40	Blood 2	1.76
	Blood 3	0.76
	Blood 4	0.18
	Blood 5	1.41
	Blood 6	1.54
45	Blood 7	0.48
	Blood 8	1.92
	Blood 9	1.63

- 39 -

Blood 10	1.65
Blood 11	1.83
Blood 12	0.37

The relative levels of expression in Table 8 show that BCSG-3 mRNA expression is detected in the pool of normal mammary gland with the highest expression value.

The absolute numbers in Table 8 were obtained analyzing pools of samples of a particular tissue from different individuals, except for the blood samples. They can not be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 9.

The absolute numbers depicted in Table 9 are relative levels of expression of BCSG-3 in 68 pairs of matching samples. All the values are compared to normal prostate (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual. In addition, 1 unmatched cancer sample (from ovary) and 1 unmatched normal sample (from ovary) were also tested.

Table 9: Relative Levels of BCSG-3 Expression in Individual Samples

Sample ID	Tissue	Cancer	Matching Normal Adjacent	Normal
Mam497M	Mammary Gland 1	3.22	1.11	
Mam173M	Mammary Gland 2	1.39	17.75	
Mam726M	Mammary Gland 3	7.62	1.31	
MamS516	Mammary Gland 4	11.08	0.10	
MamS621	Mammary Gland 5	18.25	0.05	
MamS079	Mammary Gland 6	0.78	0.24	
Mam19DN	Mammary Gland 7	71.01	1.39	
Mam522	Mammary Gland 8	3.35	0.16	

- 40 -

	MamS127	Mammary Gland 9	48.00	0.48	
	Mam162X	Mammary Gland 10	0.18	0.65	
	MamS123	Mammary Gland 11	49.69	0.00	
	MamS997	Mammary Gland 12	141.53	0.48	
5	Mam543M	Mammary Gland 13	34.66	0.10	
	Mam976M	Mammary Gland 14	0.37	0.10	
	Mam74DN	Mammary Gland 15	35.14	4.36	
	MamS918	Mammary Gland 16	16.74	5.58	
	MamS854	Mammary Gland 17	1.11	1.58	
10	Mam986	Mammary Gland 18	0.58	1.14	
	MamS967	Mammary Gland 19	121.94	2.97	
	Mam355	Mammary Gland 20	11.35	0.06	
	MamA06X	Mammary Gland 21	7.65	0.13	
	Bld32XK	Bladder 1	0.17	0.02	
15	Bld66X	Bladder 2	0.17	0.13	
	BldTR17	Bladder 3	6.21	0.00	
	Bld46XK	Bladder 4	0.06	0.00	
	BldTR14	Bladder 5	0.79	0.19	
	ClnB56	Colon 1	0.12	0.10	
20	ClnDC63	Colon 2	0.21	1.09	
	CvxKS52	Cervix 1	10.74	2.21	
	CvxNK24	Cervix 2	6.96	4.63	
	CvxKS83	Cervix 3	2.29	2.23	
	CvxNK23	Cervix 4	0.22	1.54	
25	End10479	Endometrium 1	4.68	5.13	
	End12XA	Endometrium 2	1.68	2.00	
	End5XA	Endometrium 3	0.38	0.40	
	End65RA	Endometrium 4	0.49	0.38	
	End28XA	Endometrium 5	4.32	2.94	
30	End3AX	Endometrium 6	0.21	0.21	

- 41 -

	Kid6XD	Kidney 1	0.06	0.16	
	Kid710K	Kidney 2	0.03	0.10	
	Liv175L	Liver 1	1.24	0.09	
	Liv187L	Liver 2	0.07	0.06	
5	Liv15XA	Liver 3	0.02	0.01	
	Lng47XQ	Lung 1	0.15	0.06	
	LngAC88	Lung 2	1.78	0.95	
	LngAC90	Lung 3	0.46	0.00	
	LngSQ80	Lung 4	1.91	0.35	
10	Ovr103X	Ovary 1	25.63	2.52	
	OvrA084	Ovary 2	7.70	3.19	
	OvrG010	Ovary 3	0.62	3.40	
	OvrG021	Ovary 4	0.09	0.45	
	Ovr1118	Ovary 5	0.13		
15	Ovr32RA	Ovary 6			2.81
	Pan77X	Pancreas 1	0.56	0.19	
	Pan82XP	Pancreas 2	0.62	0.73	
	Pro109XB	Prostate 1	0.00	0.10	
	Pro125XB	Prostate 2	0.05	0.01	
20	Skn248S	Skin 1	0.94	0.02	
	Skn287S	Skin 2	0.36	0.05	
	SmIntH89	Small Intestine 1	0.12	0.04	
	SmInt 21XA	Small Intestine 2	0.29	0.01	
25	Sto115S	Stomach 1	1.17	0.44	
	Sto15S	Stomach 2	0.15	0.18	
	StoMT54	Stomach 3	0.12	0.18	
	Thr590D	Thyroid	3.46	3.33	
	Tst647T	Testis	1.06	0.24	
30	Utr141XO	Uterus 1	2.86	0.51	
	Utr23XU	Uterus 2	0.60	0.13	

- 42 -

Utr85XU	Uterus 3	12.21	1.43	
Utr135XO	Uterus 4	2.98	3.93	

0.00= Negative

Table 9 represents 138 samples in 17 different
5 tissues. Table 8 and Table 9 represent a combined total of
174 samples in 26 human tissue types.

Comparisons of the level of mRNA expression in breast
cancer samples and the normal adjacent tissue from the same
individuals are shown in Table 8. BSCG-3 is expressed at
10 higher levels in 17 of 21 (81%) cancer samples (mammary
gland 1, 3-6, 7-9, 11-16, 19-21) compared to normal
adjacent tissue.

- 43 -

What is claimed is:

1. A BCSG comprising:
 - (a) a polynucleotide of SEQ ID NO:1, 2, 3, 4, 5, 18 or 20 or a variant thereof;
 - 5 (b) a protein expressed by a polynucleotide of SEQ ID NO:1, 2, 3, 4, 5, 18 or 20 or a variant thereof; or
 - (c) a polynucleotide which is capable of hybridizing under stringent conditions to the antisense sequence of SEQ ID NO: 1, 2, 3, 4, 5, 18 or 20.
- 10 2. The BCSG of claim 1 comprising a protein of SEQ ID NO:19 or 21.
3. A method for diagnosing the presence of breast cancer in a patient comprising:
 - (a) determining levels of BCSG in cells, tissues or
 - 15 bodily fluids in a patient; and
 - (b) comparing the determined levels of BCSG with levels of BCSG in cells, tissues or bodily fluids from a normal human control, wherein a change in determined levels of BCSG in said patient versus normal human control is
 - 20 associated with the presence of breast cancer.
4. A method of diagnosing metastases of breast cancer in a patient comprising:
 - (a) identifying a patient having breast cancer that is not known to have metastasized;
 - 25 (b) determining BCSG levels in cells, tissues, or bodily fluid from said patient; and
 - (c) comparing the determined BCSG levels with levels of BCSG in cells, tissue, or bodily fluid of a normal human control, wherein an increase in determined BCSG levels in
 - 30 the patient versus the normal human control is associated with breast cancer which has metastasized.

- 44 -

5. A method of staging breast cancer in a patient having breast cancer comprising:

- (a) identifying a patient having breast cancer;
- (b) determining BCSG levels in a sample of cells,
5 tissue, or bodily fluid from said patient; and
- (c) comparing determined BCSG levels with levels of BCSG in cells, tissues, or bodily fluid of a normal human control, wherein an increase in determined BCSG levels in
10 said patient versus the normal human control is associated with breast cancer which is progressing and a decrease in the determined BCSG levels is associated with breast cancer which is regressing or in remission.

6. A method of monitoring breast cancer in a patient for the onset of metastasis comprising:

- 15 (a) identifying a patient having breast cancer that is not known to have metastasized;
- (b) periodically determining levels of BCSG in samples of cells, tissues, or bodily fluid from said patient; and
- (c) comparing the periodically determined BCSG levels
20 with levels of BCSG in cells, tissues, or bodily fluid of a normal human control, wherein an increase in any one of the periodically determined BCSG levels in the patient versus the normal human control is associated with breast cancer which has metastasized.

25 7. A method of monitoring a change in stage of breast cancer in a patient comprising:

- (a) identifying a patient having breast cancer;
- (b) periodically determining levels of BCSG in cells,
tissues, or bodily fluid from said patient; and
- 30 (c) comparing the periodically determined BCSG levels with levels of BCSG in cells, tissues, or bodily fluid of a normal human control, wherein an increase in any one of the periodically determined BCSG levels in the patient versus

- 45 -

the normal human control is associated with breast cancer which is progressing in stage and a decrease is associated with breast cancer which is regressing in stage or in remission.

5 8. A method of identifying potential therapeutic agents for use in imaging and treating breast cancer comprising screening molecules for an ability to bind to BCSG wherein the ability of a molecule to bind to BCSG is indicative of the molecule being useful in imaging and
10 treating breast cancer.

9. An antibody which specifically binds BCSG.

10. The antibody of claim 9 wherein the BCSG comprises SEQ ID NO: 1, 2, 3, 4, 5, 18, 19, 20 or 21.

15 11. A method of imaging breast cancer in a patient comprising administering to the patient the antibody of claim 9.

12. The method of claim 11 wherein said antibody is labeled with paramagnetic ions or a radioisotope.

20 13. A method of treating breast cancer in a patient comprising administering to the patient the antibody of claim 9.

14. The method of claim 13 wherein the antibody is conjugated to a cytotoxic agent.

25 15. A method of treating breast cancer in a patient comprising administering to the patient a molecule which downregulates expression or activity of a BCSG.

- 46 -

16. A method of inducing an immune response against a target cell expressing a BCSG comprising delivering to a human patient an immunogenically stimulatory amount of a BCSG protein so that an immune response is mounted against
5 the target cell.

17. A vaccine for treating breast cancer comprising a BCSG.

SEQUENCE LISTING

<110> Salceda, Susana
Cafferkey, Robert
Recipon, Herve
Sun, Yongming
diaDexus, Inc.

<120> A NOVEL METHOD OF DIAGNOSING, MONITORING, STAGING,
IMAGING AND TREATING BREAST CANCER

<130> DEX-0116

<140>

<141>

<150> 60/166,973

<151> 1999-11-23

<160> 21

<170> PatentIn Ver. 2.0

<210> 1

<211> 1078

<212> DNA

<213> Homo sapiens

<400> 1

```

ggatgataca agagccaaga agggacattt gagttgtgtc gcttagatag gaaagggatc 60
cagggaaaat caacagtaag tgaggatgag cagcgtctct tggttttcat tgaggataga 120
gtaagagatt gagtttagat tgcaacagaa ggaattagtt tagataccag gaagaacttc 180
ctagcctgaa gatttgtcat agtgtctgct ttctagatat ctgggaaaga tttgataata 240
gttgtttgtg aatagaaaag aggatatgat gtttttattg gccatthttgc gggactcttc 300
gacttcttgc tgctgtctct tgaggataga ttccaattcc atcctggcga gatccaagtg 360
cttacgtact gtctccttag ctgccttaga gtaaacgac atcagttcaa tggacaaaaa 420
tcaccttcag ccatgtggtt tcttcatcat catggatttc ttttggttga caaacattct 480
ggctctcaga tgcaaaaagt cacactggga aatgaactgt aagtggtgaa attagttttg 540
gtatttaatt taaaactaca ttttagtttt tctcttctct tctatgttgc aatgaatgta 600
aagtatttgg gatccagtgc ttataaacct ttccttcctt tgtgcacaga atgtaactag 660
caagcccatt agcaccaga taattctatc atggttagttt cccatcctgg aaaatctttg 720
tacagtggga agttccccga tgtgtttttc tttcttaggt gaagggttgg ctatatcact 780
ttattgaatt ttgcattcct tagactttta aaatatacta atgtattcta gtcttactct 840
aaagaccttt gatgttaaag gaatccttca tttatttcat attccctatc tcatagggcc 900
acaattattt taatacagag atgattttca aaatatttta acaactggta caggacagat 960
gccagccact cagaagggat gcctgctgta aacaagcagt atgtatgggt gtaccaatgc 1020
ctattggctg aacattatgc tactttcaga tattaataatg gtgttccttt gaatcgtg 1078

```


<210> 2
 <211> 1713
 <212> DNA
 <213> Homo sapiens

<400> 2
 atcgcattgc accaggatga ctctgaaatg gacttcagtt cttctgctga tacatctcca 60
 gttgttactt tagctctggg agttgtggaa aagtgtcgtt gtgggccgca gaatacagcc 120
 attggatgaa tatgaagaca atcctgaaag agcttgttca gagaggatcat gaggtgactg 180
 tactggcatc ttcagcttcc attctttttg atcccaatga tgcattccact cttaaatttg 240
 aagtttatcc tacatcttta actaaaactg aatttgagaa tatcatcatg caacagggtta 300
 agagatgggc agacattcga aaagatagct tttggttata tttttcacia gaacaagaaa 360
 tcctgtggga attatatgac atatttagaa acttctgtaa agatgtagtt tcaaataaga 420
 aagttatgaa aaaactacaa gagtcaagat ttgacatcgt ttttgcatg gctgtttttc 480
 cctgtgggta gctgctggct gcgctactta acatacgggt tgtgtacagt ctccgcttta 540
 ctcttggtca cacaattgaa aggcacagtg gaggactgat tttccctcct tctacatac 600
 ctattgttat gtcaaaatta agtgatcaaa tgactttcat ggagagggtta aaaaatatga 660
 tctatgtgct ttattttgac ttttggttcc aaatgtctga tatgaagaag tgggatcagt 720
 ttacagtga agtttttaga agaccacta cttatttga gacaatggga aaagctgaca 780
 tatggcttat gcgaaactcc tggagttttc aatttctca tccattctta ccaaactgtg 840
 attttgttg aggattccac tggcaaactc gccaaacccc tacctaagga aatggaggag 900
 tttgtacaga gctctggaga aaatggtgtt gtggtgtttt ctctgggggc agtgataagt 960
 aacatgacag cagaaaaggc caatgtaatt gcaacagccc ttgccaagat cccacaaaag 1020
 gttctgtgga gatttgatgg gaataaacca gatgccttag gtctcaatac tcggctgtat 1080
 aagtggatac cccagaatga cttcttaggt catccaaaaa ccagagcttt tataactcat 1140
 ggtggagcca atggcatcta tgaggcaatc taccatggga tccctatggt gggcattcca 1200
 ttgttttggg atcaacctga taacattgct cacatgaagg ccaaggaggc agctgttaga 1260
 ttggacttca acacaatgtc gactacagac ctgctgaatg cactgaagac agtaattaat 1320
 gatcctttat ataaagagaa tattatgaaa ttatcaagaa ttcaacatga tcaaccagta 1380
 aagcccctgg atcgagcagt cttctggatt gaatttgtca tgccccacaa aggagccaaa 1440
 caccttcgag ttgcagccca tgacctcacc tgggtccagt accactcttt ggatgtgatt 1500
 ggggtttctgc tggcctgtgt ggcaactgtg atatttatca tcacaaagtt ttgtctgttt 1560
 tgtttctgga agtttgctag aaaagggaag aagggaagaa gagattagtt atgtctgaca 1620
 tttgaagctg gaaaaccaga tagataggac aacttcagtt tattccagca agaaagaaaa 1680
 gattgttatg caagatttct ttcttctgtg gac 1713

<210> 3
 <211> 2327
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (924)

<220>
 <221> unsure
 <222> (1678) .. (1704)

<400> 3

```

gatggatgca tctcaaaatg tatagccaga cttgagaggt gacaattaaa gatctaaaaa 60
agagaggaga ttcccccaaa caacaatatt taattttctt agtaaaaaga ataacagaat 120
gcatcgtggc aatccttaag caacattatc tatgtggact gcttaaatca gcaaaacacc 180
agaagtttgg ttaacttggg caatatgaca agtattactt tttgggcaaa actactcatt 240
aagcaatttc tctagtgtgt cggacacaaa taggttcttt atttttggca tgtatgcctt 300
tttattttca ttcaattttt tttttttctc agacagacat agtagtaacg actagcattg 360
gaaaatacat atcactattc ttggaatatt tatggtcagt ctacttttta gtagaatatt 420
tttggatagc gttgacacga tagatcttat tccatacttc tttattattg ataattttat 480
tttcattttt tgctttcatt attatacata ttttggtgga gaagaggttg ggcttttttg 540
aaagagacaa aaattttatta taacactaaa cactcctttt ttgacatatt aaagccttta 600
ttccatctct caagatatat tataaaattt atttttttta ttttaagattt ctgaattatt 660
ttatcttaaa ttgtgatttt aaacgagcta ttatggtacg gaactttttt taatgaggaa 720
tttcatgatg atttaggaat tttctctctt ggaaaaggct tcccctgtga tgaaaatgat 780
gtgccagcta aaattgtgtg ccatttaaaa actgaaaata ttttaaaatt atttgtctat 840
attctaaatt gagctttgga tcaaaactta ggccaggacc agctcatgcg ttctcattct 900
tccttttctc actctttctc tcancactca cctctgtatt cattctgttg tttgggatag 960
aaaaatcata aagagccaac ccactcaga acgttgtgga ttgagagaga cactacatga 1020
ctccaagtat atgagaaaag gacagagctc taattgataa ctctgtagt caaaaggaaa 1080
agagtatgcc caattctctc tacatgacat attgagattt tttttaatca acttttaaga 1140
tagtgatgtt ctgttctaaa ctgttctgtt ttagtgaagg tagattttta taaaacaagc 1200
atggggattc ttttctaagg taatattaat gagaaggga aaaagtatct ttaacagctc 1260
tttggtgaag cctgtggtag cacattatgt ttataattgc acatgtgcac ataatctatt 1320
atgatccaat gcaaatacag ctccaaaaat attaaatgta tatatatttt aaaatgcctg 1380
aggaaataca tttttcttaa taaactgaag agtctcagta tggctattaa aataattatt 1440
agcctcctgt tgtgtggctg caaaacatca caaagtgacc ggtcttgaga cctgtgaact 1500
gctgccctgt ttagtaaata aaattaatgc atttctagag ggggaatatc tgccatccag 1560
tggtggaaat gtggagtaaa gaagctggtg gtctgcttct gtgctgtatg ccagcctttt 1620
gccttaagtt gagaggaggt caactttagc tactgtcttt ggtttgagag ccatggcnnn 1680
nnnnnnnnnn nnnnnnnnnn nnnngtcgtc tttggtgagc cagtaagggtg aaagcttgct 1740
gactgtccaa ggcacaagag aaaattgagg aattgaaatg caacctgagt atcaaaactaa 1800
atattctaata caaaggtagg tactgttagg tggaattcta tcagcaggca actgcaaag 1860
agaagaagat agaaggacgc ccgtcgggac tttggagggc agtggtattt tcccaaagaa 1920
agacggccaa gggcagaggc atggattctt tgcagagcac ttccttttgg tttttcagta 1980
ctgtttcata gacagtgggc tcacatgttc ctgatagtgc tgcagttgct tagaaagcat 2040
cccagttatt gcagtaatta gaacttctgg aatatgctag ggcagaagta tgtcaagtat 2100
gtcacatgaa gaaaatgtga aattcaagag taatccacac gtgagaaaact agacaatgta 2160
cattcatgtg ttctcttgaa aggaaaggga gagctgtaag cttcactctg tctacaccg 2220
gagaaaagca ggaataactt taccgtggaa ataatgttta gcttttatgc agagaaaatt 2280
gtccttctta gagcatagag tcccaaaact caattctggt tttccc 2327

```

<210> 4

<211> 639

<212> DNA

<213> Homo sapiens

<400> 4

```

ccagaaccga gtttaggtcc aggttctcgt tctggcaaat ctttctcctt accttcttcc 60
tccacccctc cacctatgcc atgttttccc ttagccactc cccagctcgg tggaggaaaag 120
gcaggcctaa ctaggtaccg tcttcccgac tttgctcaat gatagctggg tgggtctagc 180
tgggttccag ccacttgtaa tgtgggacat ctctcaccac aactttgtag gtggagcaac 240
tgctacagag gtaaatatga ttaactttac attccatctt tcgtctgctc ccaaacttaa 300
cagcaggtaa tctgcttcta gcaagtgggtg aaggtaagag aagcatctgt ataggaggca 360
agagatctga gtccttttga aggcctatcc tctgctctgt atctcaatta ctgttcttca 420
tttcaattat tcttacctac tattcagttc ccttgatctt ttcttcttgg gggctgtctt 480
agggtcaggg agattgcaga agcaccagaa ctaggagcag ccctgagaca tggggagttg 540
gagctgaagg aggaatggca ggatgaagaa ttccctaggt gaggacgtgt gagggtggtc 600
gggagaaggg aggggtgggc acgaatggac ggaggggat 639

```

<210> 5

<211> 779

<212> DNA

<213> Homo sapiens

<400> 5

```

gtatacatc tttattaatc attttgcttc caacccatt tagcctgcc ttgaaatgca 60
aaagtctgtt ccaaataaag ccttgaatt gaagaatgaa caaacattga gagcagatga 120
gatactccca tcagaatcca aacaaaagga ctatgaagaa agttcttggg attctgagag 180
tctctgtgag actgtttcac agaaggatgt gtgtttacc aaggctacac atcaaaaaga 240
aatagataaa ataaatggaa aattagaagg gtctcctgtt aaagatggtc ttctgaaggc 300
taactgcgga atgaaagttt ctattccaac taaagcctta gaattgatgg acatgcaaac 360
tttcaaagca gagcctcccg agaagccatc tgccttcgag cctgccattg aaatgcaaaa 420
gtctgttcca aataaagcct tggaattgaa gaatgaacaa acattgagag cagatcagat 480
gttcccttca gaatcaaac aaaagaagg tgaagaaaat tcttgggatt ctgagagtct 540
ccgtgagact gtttcacaga aggatgtgtg tgtaccacag gctacacatc aaaaagaaat 600
ggataaaata agtggaaaat tagaagattc aactagccta tcaaaaatct tggatacagt 660
tcattcttgt gaaagagcaa gggaacttca aaaagatcac tgtgaacaac gtacaggaaa 720
aatggaacaa atgaaaaaga agttttgtgt actgaaaaag aaactgtcag aagcaaaaa 779

```

<210> 6

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 6

```

accccattha gcctgccat

```

19

<210> 7

<211> 24

<212> DNA

<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic

<400> 7
atgggagtat ctcatctgct ctca 24

<210> 8
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic

<400> 8
tgtttggttca ttcttcaatt ccaaggcttt 30

<210> 9
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic

<400> 9
gcccattagc acccagataa t 21

<210> 10
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic

<400> 10
gccaaccctt cacctaagaa a 21

<210> 11
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic

<400> 11

cttcccactg tacaaagatt ttccaggatg 30

<210> 12

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 12

cctggagttt tcaatttcct ca 22

<210> 13

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 13

ccccagagaa aacaccacaa 20

<210> 14

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 14

actcctccat ttccttaggt aggggtttg 29

<210> 15

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 15

gagcacttcc ttttggtttt tc 22

<210> 16

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 16

gccctagcat attccagaag ttc

23

<210> 17

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 17

tagacagtgg gctcacatgt tcctgatagt g

31

<210> 18

<211> 1722

<212> DNA

<213> Homo sapiens

<400> 18

tgcaccagga tgactctgaa atggacttca gttcttctgc tgatacatct cagttgttac 60
 ttttagctctg ggagttgtgg aaaagtgtcg gtgtgggccg cagaatacag ccattggatg 120
 aatatgaaga caatcctgaa agagcttggt cagagaggtc atgagggtgac tgtactggca 180
 tcttcagctt ccattctttt tgatcccaat gatgcatcca ctcttaaatt tgaagtttat 240
 cctacatctt taactaaaac tgaatttgag aatatcatca tgcaacaggt taagagatgg 300
 tcagacattc gaaaagatag cttttggtta ttttttcac aagaacaaga aatcctgtgg 360
 gaattatatg acatatttag aaacttctgt aaagatgtag tttcaaataa gaaagttatg 420
 aaaaaactac aagagtcaag atttgacatc gtttttgcag atgctgtttt tccctgtggg 480
 gagctgctgg ctgcgtact taacatacgg tttgtgtaca gtctccgctt tactcctggc 540
 tacacaattg aaaggcacag tggaggactg attttccctc ctctctacat acctattgtt 600
 atgtcaaaat taagtgatca aatgactttc atggagaggg taaaaaatat gatctatgtg 660
 ctttattttg acttttggtt ccaaattgtc gatatgaaga agtgggatca gttttacagt 720
 gaagtttttag gaagaccac taccttattt gagacaatgg gaaaagctga catatggctt 780
 atgcgaaact cctggagttt tcaatttctt catccattct taccaaacgt tgattttgtt 840
 ggaggattcc actgcaaacc tgccaaaccc ctacctagg aaatggagga gtttgtacag 900
 agctctggag aaaatggtgt tgtggtgttt tctctgggt cagtataag taacatgaca 960
 gcagaaaggg ccaatgtaat tgcaacagcc cttgccaaga tcccacaaa ggttctgtgg 1020
 agatttgacg ggaataaacc agatgcctta ggtctcaata ctcggtgta caagtggata 1080
 cccagaaatg accttctagg tcatccaaa accagagctt ttataactca tgggtggagcc 1140
 aatggcatct atgaggcaat ctaccatggg atccctatgg tgggcattcc attgtttttt 1200
 gatcaacctg ataacattgc tcacatgaag gccaaaggag cagctgttag attggacttc 1260
 aacacaatgt cgagtacaga cctgctgaat gcactgaaga cagtaattaa tgatccttta 1320

tataaagaga atattatgaa attatcaaga attcaacatg atcaaccagt aaagcccctg 1380
gatcgagcag tcttctggat tgaatttgtc atgccccaca aaggagccaa acaccttcga 1440
gttgagcccc atgacctcac ctgggtccag taccactctt tggatgtgat tgggtttctg 1500
ctggcctgtg tggcaactgt gatatttata atcacaaagt tttgtctgtt ttgtttctgg 1560
aagtttgcta gaaaagggaa gaagggaaaa agagattagt tatgtctgac atttgaagct 1620
ggaaaaccag atagatagga caacttcagt ttattccagc aagaaagaaa agattgttat 1680
gcaagatttc tttcttctg tgacaaaaaa aaaaaaaaaa aa 1722

<210> 19

<211> 529

<212> PRT

<213> Homo sapiens

<400> 19

Met Thr Leu Lys Trp Thr Ser Val Leu Leu Leu Ile His Leu Ser Cys
1 5 10 15

Tyr Phe Ser Ser Gly Ser Cys Gly Lys Val Leu Val Trp Ala Ala Glu
20 25 30

Tyr Ser His Trp Met Asn Met Lys Thr Ile Leu Lys Glu Leu Val Gln
35 40 45

Arg Gly His Glu Val Thr Val Leu Ala Ser Ser Ala Ser Ile Leu Phe
50 55 60

Asp Pro Asn Asp Ala Ser Thr Leu Lys Phe Glu Val Tyr Pro Thr Ser
65 70 75 80

Leu Thr Lys Thr Glu Phe Glu Asn Ile Ile Met Gln Gln Val Lys Arg
85 90 95

Trp Ser Asp Ile Arg Lys Asp Ser Phe Trp Leu Tyr Phe Ser Gln Glu
100 105 110

Gln Glu Ile Leu Trp Glu Leu Tyr Asp Ile Phe Arg Asn Phe Cys Lys
115 120 125

Asp Val Val Ser Asn Lys Lys Val Met Lys Lys Leu Gln Glu Ser Arg
130 135 140

Phe Asp Ile Val Phe Ala Asp Ala Val Phe Pro Cys Gly Glu Leu Leu
145 150 155 160

Ala Ala Leu Leu Asn Ile Arg Phe Val Tyr Ser Leu Arg Phe Thr Pro
165 170 175

Gly Tyr Thr Ile Glu Arg His Ser Gly Gly Leu Ile Phe Pro Pro Ser

180					185					190					
Tyr	Ile	Pro	Ile	Val	Met	Ser	Lys	Leu	Ser	Asp	Gln	Met	Thr	Phe	Met
195					200					205					
Glu	Arg	Val	Lys	Asn	Met	Ile	Tyr	Val	Leu	Tyr	Phe	Asp	Phe	Trp	Phe
210					215					220					
Gln	Met	Ser	Asp	Met	Lys	Lys	Trp	Asp	Gln	Phe	Tyr	Ser	Glu	Val	Leu
225					230					235					
Gly	Arg	Pro	Thr	Thr	Leu	Phe	Glu	Thr	Met	Gly	Lys	Ala	Asp	Ile	Trp
245					250					255					
Leu	Met	Arg	Asn	Ser	Trp	Ser	Phe	Gln	Phe	Pro	His	Pro	Phe	Leu	Pro
260					265					270					
Asn	Val	Asp	Phe	Val	Gly	Gly	Phe	His	Cys	Lys	Pro	Ala	Lys	Pro	Leu
275					280					285					
Pro	Lys	Glu	Met	Glu	Glu	Phe	Val	Gln	Ser	Ser	Gly	Glu	Asn	Gly	Val
290					295					300					
Val	Val	Phe	Ser	Leu	Gly	Ser	Val	Ile	Ser	Asn	Met	Thr	Ala	Glu	Arg
305					310					315					
Ala	Asn	Val	Ile	Ala	Thr	Ala	Leu	Ala	Lys	Ile	Pro	Gln	Lys	Val	Leu
325					330					335					
Trp	Arg	Phe	Asp	Gly	Asn	Lys	Pro	Asp	Ala	Leu	Gly	Leu	Asn	Thr	Arg
340					345					350					
Leu	Tyr	Lys	Trp	Ile	Pro	Gln	Asn	Asp	Leu	Leu	Gly	His	Pro	Lys	Thr
355					360					365					
Arg	Ala	Phe	Ile	Thr	His	Gly	Gly	Ala	Asn	Gly	Ile	Tyr	Glu	Ala	Ile
370					375					380					
Tyr	His	Gly	Ile	Pro	Met	Val	Gly	Ile	Pro	Leu	Phe	Phe	Asp	Gln	Pro
385					390					395					
Asp	Asn	Ile	Ala	His	Met	Lys	Ala	Lys	Gly	Ala	Ala	Val	Arg	Leu	Asp
405					410					415					
Phe	Asn	Thr	Met	Ser	Ser	Thr	Asp	Leu	Leu	Asn	Ala	Leu	Lys	Thr	Val
420					425					430					
Ile	Asn	Asp	Pro	Leu	Tyr	Lys	Glu	Asn	Ile	Met	Lys	Leu	Ser	Arg	Ile

435	440	445
Gln His Asp Gln Pro Val Lys Pro Leu Asp Arg Ala Val Phe Trp Ile		
450	455	460
Glu Phe Val Met Pro His Lys Gly Ala Lys His Leu Arg Val Ala Ala		
465	470	475 480
His Asp Leu Thr Trp Phe Gln Tyr His Ser Leu Asp Val Ile Gly Phe		
485	490	495
Leu Leu Ala Cys Val Ala Thr Val Ile Phe Ile Ile Thr Lys Phe Cys		
500	505	510
Leu Phe Cys Phe Trp Lys Phe Ala Arg Lys Gly Lys Lys Gly Lys Arg		
515	520	525

Asp

<210> 20
 <211> 10006
 <212> DNA
 <213> Homo sapiens

<400> 20
 ttcctccgcg aaggctcctt tgatattaat agtgttggtg tcttgaaact gacgtaatgc 60
 gcggagactg aggtcctgac aagcgataac atttctgata aagaccgat cttactgcaa 120
 tctctagcgt cctctttttt ggtgctgctg gtttctccag acctcgcgtc ctctcgattg 180
 ctctctcgcc ttcctatttc tttttttttt ttttaaacia aaaacaacac cccctcccct 240
 ctcccacccg gcaccgggca catccttget ctatttcctt tctctttctc tctctctctc 300
 tctctttttt aataaggggtg ggggaggga agggggggga ggcaggaaag acctttttct 360
 ctcccccccg caataatcca agatcaactc tgcaaacac agaagacggg tcatggcttt 420
 ggccgcgcgc ccaccatctt tcgggctgcc gaggggtgtt ttgacgatta atcaacagat 480
 gtacagatca gctctcaaaa tgtcttctgt gtcttctgag cgtcttctaa gacaattgca 540
 ttagcctcct gctagttgac taatagaatt aataattgta aaaagcactc taaagccaca 600
 tgccttatga agtcaatgct gggtatgatt ttacaaatat ggtccgaaa aagaaccccc 660
 ctctgagaaa cgttgcaagt gaaggcgagg gccagatcct ggagcctata ggtacagaaa 720
 gcaaggatc tggaagaac aaagaattct ctgcagatca gatgtcagaa aatacggatc 780
 agagtgatgc tgcagaacta aatcataagg aggaacatag cttgcatgtt caagatccat 840
 cttctagcag taagaaggac ttgaaaagcg cagttctgag tgagaaggct ggcttcaatt 900
 atgaaagccc cagtaaggga ggaaactttc cctcctttcc gcatgatgag gtgacagaca 960
 gaaatatgtt ggctttctca tttccagctg ctgggggagt ctgtgagccc ttgaagtctc 1020
 cgaaagagc agaggcagat gaccctcaag atatggcctg caccctca ggggactcac 1080
 tggagacaaa ggaagatcag aagatgtcac caaaggctac agaggaaaca gggcaagcac 1140
 agagtggatc agccaattgt caaggtttga gccagtttc agtggcctca aaaaaccac 1200
 aagtgccttc agatgggggt gtaagactga ataaatccaa aactgactta ctggtgaatg 1260

```

acaaccaga cccggcacct ctgtctccag agcttcagga ctttaaatgc aatatctgtg 1320
gatatggtta ctacggcaac gacccacacag atctgattaa gcacttccga aagtatcact 1380
taggactgca taaccgcacc aggcaagatg ctgagctgga cagcaaaatc ttggcccttc 1440
ataacatggg gcagttcagc cattccaaag acttccagaa ggtcaaccgt tctgtgtttt 1500
ctgggtgtgct gcaggacatc aattcttcaa ggctgtttt actaaatggg acctatgatg 1560
tgcaggtgac ttcaggtgga acattcattg gcattggacg gaaaacacca gattgccaa 1620
ggaacaccaa gtatttcgc tgtaaattct gcaatttcac ttatatgggc aactcatcca 1680
ccgaattaga acaacatttt cttcagactc acccaaacia aataaaagct tctctccct 1740
cctctgaggt tgcaaacct tcagagaaaa actctaacia gtccatccct gcacttcaat 1800
ccagtgattc tggagacttg ggaaaatggc aggacaagat aacagtcaaa gcaggagatg 1860
acactcctgt tgggtactca gtgccataa agccctcga ttctctaga caaaaggtac 1920
agaggccacc agttactact ggtgtaaat ttgtagtctc agctgtgagt catctagctc 1980
acttaaacctg ctagaacatt atggcaagca gcacggagca gtgcagtcag gcggccttaa 2040
tccagagtta aatgataagc ttccagggg ctctgtcatt aatcagaatg atctagccaa 2100
aagttcagaa ggagagacaa tgaccaagac agacaagagc tcgagtggg ctaaaaagaa 2160
ggacttctcc agcaaggag ccgaggataa tatggtaacg agctataatt gtcagtctg 2220
tgacttccga tattccaaaa gccatggccc tgatgtaatt gtagtggggc cacttctccg 2280
tcattatcaa cagctccata acattcacaa gtgtaccatt aaacactgtc cattctgtcc 2340
cagaggactt tgcagcccag aaaagcacct tggagaaatt acttatccgt ttgctttag 2400
aaaaagtaat tgttccact gtgcactctt gcttctgcac ttgtctctg gggcggtg 2460
aagctcgca gtcaaacatc agtgccatca gtgttcattc accaccctg acgtagatg 2520
actcctctt cactatgaaa gtgtgcatga gtccaagca tcggatgtca aacaagaagc 2580
aatcacctg caaggatcgg atgggcagca gtctgtcaag gaaagcaaag aacactcatg 2640
taccaaatgt gatattatta ccaagtggg agaagagatt tcccgacact acaggagagc 2700
acacagctgc tacaatgcc gtcagtgcag tttacagct gccgatactc agtcactact 2760
ggagcacttc aacactgttc actgccagga acaggacatc actacagcca acggcgaaga 2820
ggacggtcat gccatatcca ccatcaaga ggagccaaa attgacttca ggggtctaaa 2880
tctgctaact ccagactcta aaatgggaga gccagttct gagagtgtg tgaagagaga 2940
gaagctggaa gagaaggacg ggctcaaaga gaaagtttg accgagagt ccagtgatga 3000
ccttcgcaat gtgacttggg gaggggcaga catcctgcgg gggagtccgt catacacca 3060
agcaagcctg gggctgctga cgctgtgtc tggcaccaa gagcagacaa agactctaag 3120
ggatagtccc aatgtggagg ccgccatct ggcgacact atttatggct tggctgtgga 3180
aaccaaggga ttctgcagg gggcgccagc tggcgagag agtctgggg ccctcccca 3240
gcagtatcct gcatcgggag aaaacaagtc caaggatgaa tcccagtccc tgttacggag 3300
gcgtagaggc tccggtgtt tttgtgcaa ttgcctgacc acaaagacct ctctctggcg 3360
aaagaatgca aatggcgat atgtatgcaa cgctgtggc ctctaccaga agcttcaact 3420
gactccagg cctttaaaac tcattaaaca aaacaacggt gagcagatta ttaggaggag 3480
aacaagaaag cgccttaacc cagaggcact tcaggctgag cagctcaaca aacagcagag 3540
gggcagcaat gaggagcaag tcaatggaag ccggttagag aggaggtcag aagatcatct 3600
aactgaaagt caccagagag aaattccact cccagccta agtaaatagc aagccaggg 3660
ttcattgact aaaagccatt ctgctcagca gccagtcctg gtcagccaaa ctctggatat 3720
tcacaaaagg atgcaacct tgcacattca gataaaaagt cctcaggaaa gtactggaga 3780
tccaggaaat agttcatccg tatctgaagg gaaaggaagt tctgagagag gcagtcctat 3840
agaaaagtac atgagacctg cgaaacaccc aaattattca ccaccaggca gccctattga 3900
aaagtaccag taccacttt ttggacttcc cttttacata atgacttcca gagtgaagct 3960
gattggctgc ggttctggg taaatataag ctctccgttc ctgggaatcc gcactacttg 4020
agtcacgtgc ctggcctacc aaatccttgc caaaactatg tgccttatcc cacttcaat 4080
ctgcctctc attttctcag tgttgatca gacaatgaca ttctctaga tttggcgatc 4140

```

```

aagcattcca gacctgggcc aactgcaaac ggtgcctcca aggagaaaac gaaggcacca 4200
ccaaatgtaa aaaatgaagg tcccttgaat gtagtaaaaa cagagaaagt tgatagaagt 4260
actcaagatg aactttcaac aaaatgtgtg cactgtggca ttgtctttct ggatgaagtg 4320
atgtatgctt tgcataatgag ttgccatggt gacagtggac ctttccagtg cagcatatgc 4380
cagcatcttt gcacggacaa atatgacttc acaacacata tccagagggg cctgcatagg 4440
aacaatgcac aagtggaaaa aaatggaaaa cctaaagagt aaaaccttag cacttagcac 4500
aattaaatag aaatagggtt tcttgatggg aattcaatag cttgtaatgt cttatgaaga 4560
cctattaaaa aaatacttca tagagcctgc cttatccaac atgaaattcc cttcttttgt 4620
tattctttct tttgatgagt aggttaccac gattaaaaag tgagataaat ggtcaatgag 4680
aaagaatgga agatggtaaa caatcacttt ttaaacctg ttaagtcaaa accatcttgg 4740
ctaatatgta ctggggaaat aatccataag agatatcacc agactagaat taatatattt 4800
ataaagaaag agacaaaaac tgtctagaat ttgaaagggg ttacatatta ttatactaaa 4860
gcagtactgg actggccatt ggaccatttg ttccaaaacc cataaattgt tgcctaaatt 4920
tataatgatc atgaaaccct aggcagagga ggagaaattg aagggtccagg gcaatgaaag 4980
aaaaatggcg ccctctcaat ttagtcttct ctcatggcc atgtttcaga ttttgacctt 5040
gaaatgcgag ctgtgggttag gcttgggttag agtgcagcaa gcaacatgac agatgggtggc 5100
acgctgtttt taccagccc tgccgtgaca tacacatgca caccctctct gatatttttg 5160
tcctttagat gttcaaatac tcagtagtcc ttttgtttgc ggtttagatt cattttgtcc 5220
acacatgtac ccatttttaa aaacaatgtc ctcgatgctt ctgtagtgat ttcattttag 5280
ccaggatatt ctttcttgtg tgtgatgaac cagtatggat ttgcttttct aagcctcctg 5340
ttggttacta atctcacttg gcacattata actaaaggaa tcccctcaat tcaaaagcat 5400
agatggatac aaatgtcaga ccgtgggttt aatttgttta gaacacatgg catttcttca 5460
caaggtaacc tgctgtattt atttattttc ttttgggttaa atataatttc caaactttgt 5520
ggtcaggcag cgtctaaggt tacgttacca cagactgaca gttggtatat gtaccagcca 5580
atcccttcat taaatgtata cagatttagt taagtagcat taaataggat tcttagaagt 5640
atgtcctcat agaactttta atacttaagg ctttgtaaaa actatccatg aagggaaagc 5700
tcctcagcat aactgctcag ggaaataggg ctaaaataact gaacattaaa taattgggtt 5760
aagggtgctg tagtcgagcc tcaatgcttg ctacaaggat gtatgtacaa ggactgactt 5820
taataatttg cattatattg tcccaaccag tagtttattt tttgccacgg agatgtagaa 5880
gatattacaa gctactggat gcactgtcag attaaacttat ttcattaaag aagttgggag 5940
aacaatatag aaaaaaaaaa ttatttttct agtaaatatt aatgtattac atttcaaata 6000
atgggtgcctg acatattgaa taattatttt ctacagtgtc cgtatgcaac aaagatattc 6060
catcatgcat tagagtcagt tctggctctg cctagctgtt tacatttgca aatgtagcaa 6120
acaaggtaat gaagcaacta tttctattgc agtagatatc cttttgtgtg tgtgtgtgtg 6180
cattaaagtt gtaaacggta acatgaaaca aatgaaagtt cttgctataa tggatggaa 6240
aacaagaagg aaatgaaaat atttttatgc ctacttagga aaaaaagggg agcacttatt 6300
cattccaagt actttttttt ttttaatttt taagctctta actcacattg ttatgcttaa 6360
gatgataaac atatatcctc tttttattgc tttgtctatg tttcatatga aacatttcag 6420
aaattatttt gataagtgtt gctggaatct gcaacgctga tttttttttg cattctgtag 6480
tcgcatttgc actccatttt tacattaatt cgcagttgct ttgtatcatt gttttgtttg 6540
ggttttgttt ctttttcaca gtgccgggtc ttcgtttctt aaagtggat ggcaggtaga 6600
gttcaaccag ttcgtgactg ttgtagcgaa tgaagttaaa aaaatgtctt tctgatgttg 6660
tgttgtcatt ttcatttttg catttttttg tttgcatatt aaaaaagag aaaagagaaa 6720
gcaagagaca gaaatcagga ctaagtcctc tgcttcagtt tcattgttaa cgggccttat 6780
tctgatctca cctgtcgcgt agctctaata ttcacataaa ctgaaataaa gaagtggaat 6840
gaggagcttt gacattcaaa ttatgtgatg taatttatct tccttaggaa ttttgatgga 6900
tgcattctca aatgtatagc cagacttgag aggtgacaat taaagatcta aaaaagagag 6960
gagattcccc caaacaacaa tatttaattt tcttagtaaa aagaataaca gaatgcatcg 7020

```

```

tggcaatcct taagcaacat tatctatgtg gactgcttaa atcagcaaaa caccagaagt 7080
ttgggttaact tgggcaatat gacaagtatt actttttggg caaaactact cattaagcaa 7140
tttctctagt gtgtcggaca caaatagggt ctttattttt ggcatgtatg cttttttatt 7200
ttcattcaat tttttttttt tctcagacag acatagtagt atcaactagc attggaaaat 7260
acatatcact attcttggaa tatttatggg cagtctactt tttagtaaaa tattttttgga 7320
tagcgttgac acgatagatc ttattccata cttctttatt attgataatt ttattttcat 7380
tttttgcttt cattattata catatttttg tggagaagag gttgggcttt tttgaaagag 7440
acaaaaattt attataacac taaacactcc ttttttgaca tattaaagcc tttattccat 7500
ctctcaagat atattataaa atttattttt ttaatttaag atttctgaat tattttatct 7560
taaattgtga ttttaaacga gctattatgg tacggaactt tttttaatga ggaatttcat 7620
gatgatttag gaattttctc tcttggaaaa ggcttccctt gtgatgaaaa tgatgtgcca 7680
gctaaaattg tgtgccattt aaaaactgaa aatattttta aattatttgt ctatattcta 7740
aattgagctt tggatcaaac tttaggccag gaccagctca tgcgttctca ttcttctttt 7800
tctcactctt tctctcatca ctcacctctg tattcattct gttgtttggg atagaaaaat 7860
cataaagagc caaccatct cagaacgttg tggattgaga gagacactac atgactccaa 7920
gtatatgaga aaaggacaga gctctaattg ataactctgt agttcaaaag gaaagagtat 7980
gcccaattct ctctacatga catattgaga ttttttttaa tcaactttta agatagtgat 8040
gttctgttct aaactgttct gtttttagtga aggtagattt ttataaaaca agcatgggga 8100
ttcttttcta aggtaatat aatgagaagg gaaaaaagta tctttaacag ctctttgttg 8160
aagcctgtgg tagcacatta tgtttataat tgcacatgtg cacataatct attatgatcc 8220
aatgcaaata cagctccaaa aatattaaat gtatatatat tttaaaatgc ctgaggaaat 8280
acatttttct taataaactg aagagtctca gtatggctat taaaataatt attagcctcc 8340
tgttgtgtgg ctgcaaaaca tcacaaagtg accggtcttg agacctgtga actgctgccc 8400
tgttttagtaa ataaaattaa tgcatttcta gagggggaat atctgccatc cagtgggtga 8460
aatgtggagt aaagaagctg gtggtctgct tctgtgctgt atgccagcct tttgccttaa 8520
gttgagagga ggtcaacttt agctactgtc tttggtttga gagccatggc aaaaaaaaaa 8580
aaagaaaaaa agatcaagtc gtctttgggt agccagtaag gtgaaagctt gctgactgtc 8640
caaggcacaa gagaaaattg aggaattgaa atgcaacctg agtatcaaac taaatattct 8700
aatcaaaggt aggtactgtt aggtggaatt ctatcagcag gcaactgcaa atgagaagaa 8760
gatagaagga cgcccgtcgg gactttggag ggcattgtta tttcccaa gaaagacggc 8820
caagggcaga ggcatggatt ctttgcagag cacttctttt tggtttttca gtactgttc 8880
atagacagtg ggctcacatg ttctgatag tgcctcagtt gcttagaaag catcccagtt 8940
aattgcagta attagaactt ctggaatatg ctagggcaga agtatgtcaa gtatgtcaca 9000
tgaagaaaat gtgaaattca agagtaatcc acacgtgaga aactagacaa tgtacattca 9060
tgtgttctct tgaaggaaa gggagagctg taagcttcac tctgtcctac accggagaaa 9120
agcaggaata actttaccgt ggaaataatg tttagctttt atcagagaaa attgtccttc 9180
tagagcatag agtcccaaaa ctcaattctg gttttccctt gttttttttt tttttttttt 9240
tcccaacata tgaactgcag catatcactt tttctttttt tgcctcaggt tctcacctg 9300
taaaattgaa aaatatatgt attaataata ttattaataa taataatggg aatgtagtac 9360
ttgtttgtaa agcactttga gatccttggg tgaaggcac cataggagtg ccaagtatta 9420
ttatgtggcc aagggggtta tttaaactgt cagttcccaa aggccaggaa aggttgggg 9480
catttttctt aaagacgagc tgtaaatatc aactaggcag ccaatagtgt tgactatgaa 9540
gatgcaaaac tattactagg ctgataaaat catagtctt taatggctac caataaggca 9600
aatatcacia taataaacgc caaatcctt agggcggact atttgacaac cacatggaaa 9660
actttggggg aggcattgag ggggaacatc tcaaaatgcc aatgtaaaat ttaacttaca 9720
gcaatattca ccagcagaaa atgtctttca tatggaatga tttcatgttg ctaagaaaaa 9780
gaattcaatt tgtagtcctg atttgaatac tagaatgttg gctataatag ttctgttctt 9840
acaacacatg aaattttttc gttttatttt attttgttt catagtgcac gttcatttct 9900

```

actcacaaac atgttcttgg tgtatttctt atgcaaacia tcttcaggca gcaaagatgt 9960
 ctgttacatc taaacttgaa taataaagtt ttccaccagt tacaca 10006

<210> 21

<211> 1281

<212> PRT

<213> Homo sapiens

<400> 21

Met Val Arg Lys Lys Asn Pro Pro Leu Arg Asn Val Ala Ser Glu Gly
 1 5 10 15

Glu Gly Gln Ile Leu Glu Pro Ile Gly Thr Glu Ser Lys Val Ser Gly
 20 25 30

Lys Asn Lys Glu Phe Ser Ala Asp Gln Met Ser Glu Asn Thr Asp Gln
 35 40 45

Ser Asp Ala Ala Glu Leu Asn His Lys Glu Glu His Ser Leu His Val
 50 55 60

Gln Asp Pro Ser Ser Ser Ser Lys Lys Asp Leu Lys Ser Ala Val Leu
 65 70 75 80

Ser Glu Lys Ala Gly Phe Asn Tyr Glu Ser Pro Ser Lys Gly Gly Asn
 85 90 95

Phe Pro Ser Phe Pro His Asp Glu Val Thr Asp Arg Asn Met Leu Ala
 100 105 110

Phe Ser Phe Pro Ala Ala Gly Gly Val Cys Glu Pro Leu Lys Ser Pro
 115 120 125

Gln Arg Ala Glu Ala Asp Asp Pro Gln Asp Met Ala Cys Thr Pro Ser
 130 135 140

Gly Asp Ser Leu Glu Thr Lys Glu Asp Gln Lys Met Ser Pro Lys Ala
 145 150 155 160

Thr Glu Glu Thr Gly Gln Ala Gln Ser Gly Gln Ala Asn Cys Gln Gly
 165 170 175

Leu Ser Pro Val Ser Val Ala Ser Lys Asn Pro Gln Val Pro Ser Asp
 180 185 190

Gly Gly Val Arg Leu Asn Lys Ser Lys Thr Asp Leu Leu Val Asn Asp
 195 200 205

Asn Pro Asp Pro Ala Pro Leu Ser Pro Glu Leu Gln Asp Phe Lys Cys
 210 215 220

Asn Ile Cys Gly Tyr Gly Tyr Tyr Gly Asn Asp Pro Thr Asp Leu Ile
 225 230 235 240

Lys His Phe Arg Lys Tyr His Leu Gly Leu His Asn Arg Thr Arg Gln
 245 250 255

Asp Ala Glu Leu Asp Ser Lys Ile Leu Ala Leu His Asn Met Val Gln
 260 265 270

Phe Ser His Ser Lys Asp Phe Gln Lys Val Asn Arg Ser Val Phe Ser
 275 280 285

Gly Val Leu Gln Asp Ile Asn Ser Ser Arg Pro Val Leu Leu Asn Gly
 290 295 300

Thr Tyr Asp Val Gln Val Thr Ser Gly Gly Thr Phe Ile Gly Ile Gly
 305 310 315 320

Arg Lys Thr Pro Asp Cys Gln Gly Asn Thr Lys Tyr Phe Arg Cys Lys
 325 330 335

Phe Cys Asn Phe Thr Tyr Met Gly Asn Ser Ser Thr Glu Leu Glu Gln
 340 345 350

His Phe Leu Gln Thr His Pro Asn Lys Ile Lys Ala Ser Leu Pro Ser
 355 360 365

Ser Glu Val Ala Lys Pro Ser Glu Lys Asn Ser Asn Lys Ser Ile Pro
 370 375 380

Ala Leu Gln Ser Ser Asp Ser Gly Asp Leu Gly Lys Trp Gln Asp Lys
 385 390 395 400

Ile Thr Val Lys Ala Gly Asp Asp Thr Pro Val Gly Tyr Ser Val Pro
 405 410 415

Ile Lys Pro Leu Asp Ser Ser Arg Gln Asn Gly Thr Glu Ala Thr Ser
 420 425 430

Tyr Tyr Trp Cys Lys Phe Cys Ser Phe Ser Cys Glu Ser Ser Ser Ser
 435 440 445

Leu Lys Leu Leu Glu His Tyr Gly Lys Gln His Gly Ala Val Gln Ser
 450 455 460

Gly Gly Leu Asn Pro Glu Leu Asn Asp Lys Leu Ser Arg Gly Ser Val			
465	470	475	480
Ile Asn Gln Asn Asp Leu Ala Lys Ser Ser Glu Gly Glu Thr Met Thr			
	485	490	495
Lys Thr Asp Lys Ser Ser Ser Gly Ala Lys Lys Lys Asp Phe Ser Ser			
	500	505	510
Lys Gly Ala Glu Asp Asn Met Val Thr Ser Tyr Asn Cys Gln Phe Cys			
	515	520	525
Asp Phe Arg Tyr Ser Lys Ser His Gly Pro Asp Val Ile Val Val Gly			
	530	535	540
Pro Leu Leu Arg His Tyr Gln Gln Leu His Asn Ile His Lys Cys Thr			
	545	550	555
Ile Lys His Cys Pro Phe Cys Pro Arg Gly Leu Cys Ser Pro Glu Lys			
	565	570	575
His Leu Gly Glu Ile Thr Tyr Pro Phe Ala Cys Arg Lys Ser Asn Cys			
	580	585	590
Ser His Cys Ala Leu Leu Leu Leu His Leu Ser Pro Gly Ala Ala Gly			
	595	600	605
Ser Ser Arg Val Lys His Gln Cys His Gln Cys Ser Phe Thr Thr Pro			
	610	615	620
Asp Val Asp Val Leu Leu Phe His Tyr Glu Ser Val His Glu Ser Gln			
	625	630	635
Ala Ser Asp Val Lys Gln Glu Ala Asn His Leu Gln Gly Ser Asp Gly			
	645	650	655
Gln Gln Ser Val Lys Glu Ser Lys Glu His Ser Cys Thr Lys Cys Asp			
	660	665	670
Phe Ile Thr Gln Val Glu Glu Glu Ile Ser Arg His Tyr Arg Arg Ala			
	675	680	685
His Ser Cys Tyr Lys Cys Arg Gln Cys Ser Phe Thr Ala Ala Asp Thr			
	690	695	700
Gln Ser Leu Leu Glu His Phe Asn Thr Val His Cys Gln Glu Gln Asp			
	705	710	715
			720

Ile Thr Thr Ala Asn Gly Glu Glu Asp Gly His Ala Ile Ser Thr Ile	725	730	735
Lys Glu Glu Pro Lys Ile Asp Phe Arg Val Tyr Asn Leu Leu Thr Pro	740	745	750
Asp Ser Lys Met Gly Glu Pro Val Ser Glu Ser Val Val Lys Arg Glu	755	760	765
Lys Leu Glu Glu Lys Asp Gly Leu Lys Glu Lys Val Trp Thr Glu Ser	770	775	780
Ser Ser Asp Asp Leu Arg Asn Val Thr Trp Arg Gly Ala Asp Ile Leu	785	790	795
Arg Gly Ser Pro Ser Tyr Thr Gln Ala Ser Leu Gly Leu Leu Thr Pro	805	810	815
Val Ser Gly Thr Gln Glu Gln Thr Lys Thr Leu Arg Asp Ser Pro Asn	820	825	830
Val Glu Ala Ala His Leu Ala Arg Pro Ile Tyr Gly Leu Ala Val Glu	835	840	845
Thr Lys Gly Phe Leu Gln Gly Ala Pro Ala Gly Gly Glu Lys Ser Gly	850	855	860
Ala Leu Pro Gln Gln Tyr Pro Ala Ser Gly Glu Asn Lys Ser Lys Asp	865	870	875
Glu Ser Gln Ser Leu Leu Arg Arg Arg Arg Gly Ser Gly Val Phe Cys	885	890	895
Ala Asn Cys Leu Thr Thr Lys Thr Ser Leu Trp Arg Lys Asn Ala Asn	900	905	910
Gly Gly Tyr Val Cys Asn Ala Cys Gly Leu Tyr Gln Lys Leu His Ser	915	920	925
Thr Pro Arg Pro Leu Asn Ile Ile Lys Gln Asn Asn Gly Glu Gln Ile	930	935	940
Ile Arg Arg Arg Thr Arg Lys Arg Leu Asn Pro Glu Ala Leu Gln Ala	945	950	955
Glu Gln Leu Asn Lys Gln Gln Arg Gly Ser Asn Glu Glu Gln Val Asn	965	970	975

Gly Ser Pro Leu Glu Arg Arg Ser Glu Asp His Leu Thr Glu Ser His
 980 985 990

Gln Arg Glu Ile Pro Leu Pro Ser Leu Ser Lys Tyr Glu Ala Gln Gly
 995 1000 1005

Ser Leu Thr Lys Ser His Ser Ala Gln Gln Pro Val Leu Val Ser Gln
 1010 1015 1020

Thr Leu Asp Ile His Lys Arg Met Gln Pro Leu His Ile Gln Ile Lys
 1025 1030 1035 1040

Ser Pro Gln Glu Ser Thr Gly Asp Pro Gly Asn Ser Ser Ser Val Ser
 1045 1050 1055

Glu Gly Lys Gly Ser Ser Glu Arg Gly Ser Pro Ile Glu Lys Tyr Met
 1060 1065 1070

Arg Pro Ala Lys His Pro Asn Tyr Ser Pro Pro Gly Ser Pro Ile Glu
 1075 1080 1085

Lys Tyr Gln Tyr Pro Leu Phe Gly Leu Pro Phe Val His Asn Asp Phe
 1090 1095 1100

Gln Ser Glu Ala Asp Trp Leu Arg Phe Trp Ser Lys Tyr Lys Leu Ser
 1105 1110 1115 1120

Val Pro Gly Asn Pro His Tyr Leu Ser His Val Pro Gly Leu Pro Asn
 1125 1130 1135

Pro Cys Gln Asn Tyr Val Pro Tyr Pro Thr Phe Asn Leu Pro Pro His
 1140 1145 1150

Phe Ser Ala Val Gly Ser Asp Asn Asp Ile Pro Leu Asp Leu Ala Ile
 1155 1160 1165

Lys His Ser Arg Pro Gly Pro Thr Ala Asn Gly Ala Ser Lys Glu Lys
 1170 1175 1180

Thr Lys Ala Pro Pro Asn Val Lys Asn Glu Gly Pro Leu Asn Val Val
 1185 1190 1195 1200

Lys Thr Glu Lys Val Asp Arg Ser Thr Gln Asp Glu Leu Ser Thr Lys
 1205 1210 1215

Cys Val His Cys Gly Ile Val Phe Leu Asp Glu Val Met Tyr Ala Leu
 1220 1225 1230

His Met Ser Cys His Gly Asp Ser Gly Pro Phe Gln Cys Ser Ile Cys
1235 1240 1245

Gln His Leu Cys Thr Asp Lys Tyr Asp Phe Thr Thr His Ile Gln Arg
1250 1255 1260

Gly Leu His Arg Asn Asn Ala Gln Val Glu Lys Asn Gly Lys Pro Lys
1265 1270 1275 1280

Glu